

# Author Index

- Abalos A, see Lien CA
- Abderrahmane B, Omar A. An unusual cause of a double-lumen endotracheal tube obstruction (letter), 694
- Abke J, see Kochs E
- Abramovitz S, Beilin Y, Bernstein HH. Isoproterenol and epidural analgesia (letter), 1190
- Adachi T, Fukumoto M, Uetsuki N, Yasui O, Hayashi M. Suspected severe methemoglobinemia caused by topical application of an ointment containing benzocaine around the enterostomy (letter), 1190  
see Miyazaki Y  
see Shinomura T
- Adams SS. The *Titanic* is not the best analogy for enoxaparin (letter), 229
- Adsumelli RN, see Manecke GR Jr
- Afshar M, see Goldberg ME
- Agrò F, see Varrassi G
- Ahmed HE, see Ghoname EA
- Akamatsu H, see Nishina K
- Akça O  
see Fleischmann E  
see Hoerauf KH
- Akhtar S, see Raby KE
- Albani A, see Runza M
- Aldegheri G, see Casati A
- Allan L, see Cornish PB
- Alley C, see Gwartz KH
- Allison PM, see Whitten CW
- Aloe L, see Varrassi G
- Alon E, see Laviola S
- Al-Rawi P, see Gupta AK
- Alves D, see Palmer CM
- Amaki Y, see Nishiyama M
- An G, see Xue FS
- Andelinovic S, see Kapural L
- Anderson EA, see Domino KB
- Andoh T  
see Furuya R  
see Soga K  
see Watanabe I
- Andrzejowski J, see Sleigh JW
- Angel JJ, see Ramanathan J
- Antognini JF, Carstens E, Buzin V. Isoflurane depresses motoneuron excitability by a direct spinal action: an F-wave study, 681
- Aoyama K  
see Nijima K  
see Seto A
- Apfel C, see Greim C-A
- Arkiliç CF, see Fleischmann E
- Arlander E, see Österlund Å
- Arledge JA, see Sullivan LM
- Arruda J, see Rassias AJ
- Artru AA, Momota Y. Trabecular outflow facility and formation rate of aqueous humor during anesthesia with sevoflurane-nitrous oxide or sevoflurane-remifentanyl in rabbits, 781  
Propofol for lidocaine-induced seizures (letter, reply), 1193  
see Talmor D
- Asai T, Johmura S, Shingu K. Use of the laryngeal mask before tracheal intubation in a patient with a cancerous nose (letter), 1429
- Power I. Naloxone inhibits gastric emptying in the rat, 204
- Shingu K. Use of the laryngeal mask during emergence from anesthesia in a patient with an unstable neck (letter), 469
- Aşik I, see Kurtipek Ö
- Atakan Ş, see Celebioglu B
- Ateş Y, see Kurtipek Ö
- Aubert G, see Rennotte MTh
- Aversano T, see Landesburg G
- Bachmann D, see Kapral S
- Backman SB, Stein RD, Polosa C.  
Organization of the sympathetic innervation of the forelimb resistance vessels in the cat, 320
- Baele Ph, see Rennotte MTh
- Baer GA, see Puura AIE
- Bailey AG, see Valley RD
- Bailey MK, see Burt N
- Ballabio M, see Varrassi G
- Barclay DK, see Cheung AT
- Barnes RA, see Frank SM
- Barnett RA. The MGH Board Review of Anesthesiology. 5th ed. (book review), 1195
- Barranco M, see Garutti I
- Barrett PQ, see McDowell TS
- Bart RD, see Miura Y
- Bartel M, see Klein U
- Başar HT, see Kurtipek Ö
- Bass EB, see Lehmann HP
- Bastien JL. Inhaled nitric oxide and amniotic fluid embolism (letter, reply), 691
- Bavaria JE, see Cheung AT
- Baysinger CL. The High Risk Obstetrical Patient. Vol 16, No 2 of Anesthesiology Clinics of North America (book review), 471
- Beach M, see Rassias AJ
- Beattie C. Successful strategies for improving operating room efficiency at academic institutions (letter), 963
- Becker BF, see Szekely A
- Bedel MJ, see Williams EL
- Bedforth NM, see Girling KJ
- Beemer GH, see Leslie K
- Been HD, see Ubags LH
- Begou G, see Parlow JL
- Beh T, see Myles PS
- Behringer EC. Comparison of the laryngeal mask airway and cuffed oropharyngeal airway: alternative hypotheses (letter), 961
- Beilin Y, Galea M, Zahn J, Bodian CA. Epidural ropivacaine for the initiation of labor epidural analgesia: a dose finding study, 1340  
see Abramovitz S
- Bell E, see D'Ercole FJ
- Belmont MR, see Lien CA
- Ben-David B, see Gaitini LA
- Benhamou D, see Bouaziz H
- Bennett-Guerrero E, see Gan TJ
- Berendes E, see Loick HM
- Berge KH, see Kuhnert SM
- Bernard J-M, see Gentili M
- Bernheim J, see Fredman B
- Bernstein HH, see Abramovitz S
- Berry AJ. Recommended exposure limits for desflurane and isoflurane (letter), 1424
- Berry AM, see Brimacombe JC
- Berti M, see Casati A
- Bertrand M, see Eyraud D
- Bicker AA, see Manecke GR Jr
- Bickler PE, see Eilers H
- Biehl J, see Bourke DL
- Binder M, see Oczenski W
- Birnback DJ, Chestnut DH. The epidural test dose in obstetric patients: has it outlived its usefulness?, 971
- Bischoff P, see Kochs E
- Bishop AT, see Horlocker TT
- Blake D, see Scott DA
- Blakeman B, see Jellish WS
- Blomquist H, Dahlgren G. Anesthesia: Biologic Foundations (book review), 235
- Bode C, see Gries A
- Bodian CA  
see Beilin Y  
see Reich DL

- Boezeman EH, see van Dongen EP  
 Bohrer H, see Gries A  
 Boldt J, Maleck W, Koetter KP. Which countries publish in important anesthesia and critical care journals?, 1175  
   see Suttner S  
 Bone HG, see Booke M  
 Bonnet F, see Gentili M  
 Booke M, Hagemann O, Van Aken H, Erren M, Wullenweber J, Bone HG. Intraoperative autotransfusion in small children: an *in vitro* investigation to study its feasibility, 763  
 Borrenpohl S, see Norris MC  
 Botero M, Enneking FK. Reversal of prolonged unconsciousness by naloxone after an intravascular injection of a local anesthetic and clonidine, 1185  
 Bouaziz H, Okubo N, Malinovsky J-M, Benhamou D, Samii K, Mazoit JX. The age-related effects of epidural lidocaine, with and without epinephrine, on spinal cord blood flow in anesthetized rabbits, 1302  
 Boudreaux AM, see Segura-Vasi A  
 Bourke DL, Biehl J. The laryngotracheal topical anesthesia kit with capnography for difficult endotracheal intubation, 943  
 Brabant S, see Eyraud D  
 Brack A, Schäfer M. Atlas of Anesthesia, Vol II. Scientific Principles of Anesthesia (book review), 471  
 Bradford C, see Gan TJ  
 Bradley EL Jr, see Vinik HR  
 Brederlau J, see Greim C-A  
 Breivik H  
   see Klaastad Ø  
   see Lauwers M  
 Brimacombe J, see Keller C  
 Brimacombe JC, see Brimacombe JC  
 Brimacombe JR, Brimacombe JC, Berry AM. Comparison of the laryngeal mask airway and cuffed oropharyngeal airway: alternative hypotheses (letter, reply), 961  
 Brock-Utne JG, see Torralva PR  
 Brodner G, Mertes N, Van Aken H, Pogatzki E, Buerkle H, Marcus MA, Mollhoff T. Epidural analgesia with local anesthetics after abdominal surgery: earlier motor recovery with 0.2% ropivacaine than 0.175% bupivacaine, 128  
 Brodsky JB, Fitzmaurice BG, Macario A. Selecting double-lumen tubes for small patients (letter), 466  
 Broka SM, Gourdange PA, Joucken KL. Sevoflurane and desflurane confusion (letter), 1194  
 Morimont CJ, Joucken KL. Perioperative torsade de pointes ventricular tachycardia (letter), 966  
 Bromage PR. The influence of epidural needle bevel on spread of sensory blockade in the laboring parturient (letter), 962  
 Bronsther O, see Lustik SJ  
 Brown PC, see Nagi H  
 Brown RD, see Grady RE  
 Brull SJ, see Raby KE  
 Buckland M, see Scott DA  
 Bucklin BA, Tinker JH, Smith CV. Clinical dilemma: a patient with postdural puncture headache and acute leukemia, 166  
 Buerkle H, see Brodner G  
 Buffington CW, see Nyström EUM  
 Bugter MLT, see Slappendel R  
 Burgmann H, see Knapp S  
 Burke SJ, see Manninen PH  
 Burkhardt D, see Whitten CW  
 Burm AGL, see Stienstra R  
 Burt N, Haynes GR, Bailey MK. Patients with malignant osteopetrosis are at high risk of anesthetic morbidity and mortality, 1292  
 Bush AJ, see Ramanathan J  
 Buzin V, see Antognini JF  
 Byers RS, see Gwirtz KH  
 Byrick RJ. Fat embolism and neurological dysfunction (letter), 1427  
 Cahalan MK, see Lambert A-S  
 Caldwell JE, see Talke PO  
 Calès J, see Grenier B  
 Calhoun P, see Valley RD  
 Camann W. Visual analog scale scores for labor pain (letter), 1421  
 Camporesi EM, see Thomas PS  
 Campos C, Naguib SS, Chuang AZ, Lemak NA, Khalil SN. Endobronchial intubation causes an immediate increase in peak inflation pressure in pediatric patients, 268  
 Camu F, see Lauwers M  
 Cancel D  
   see Miller KM  
   see Cassidy JF Jr  
 Candela-Toha AM, Martínez-Pérez A. Transfusion requirements in orthotopic liver transplantation (letter), 692  
 Candido KD, see Khorasani A  
 Cantillo J, see Goldberg ME  
 Cantwell MF, see Politis GD  
 Caramico LA, see Kain ZN  
 Carrero E, see Suarez S  
 Carstens E, see Antognini JF  
 Casaletti E, see Casati A  
 Casati A, Fanelli G, Casaletti E, Cedrati V, Veglia F, Torri G. The target plasma concentration of propofol required to place laryngeal mask versus cuffed oropharyngeal airway, 917  
 Fanelli G, Cedrati V, Berti M, Aldegheri G, Torri G. Pulmonary function changes after interscalene brachial plexus anesthesia with 0.5% and 0.75% ropivacaine: a double-blinded comparison with 2% mepivacaine, 587  
   see Fanelli G  
 Cassidy JF Jr, Wysocki TT, Miller KM, Cancel DD, Izenberg N. Use of a preanesthetic video for facilitation of parental education and anxiety reduction before pediatric ambulatory surgery, 246  
   see Miller KM  
 Cassuto J, see Mattsson U  
 Cedrati V, see Casati A  
 Celebioglu B, Eslambouli NR, Olcay E, Atakan S. The effect of tenoxicam on intraperitoneal adhesions and prostaglandin E<sub>2</sub> levels in mice, 939  
 Chan V, see Mulroy MF  
 Chaney MA, Nikolov MP. The relationship between dynamic compliance and inspiratory flow (letter, reply), 465  
   Port-Access Cardiac Surgery for the Anesthesiology (book review), 235  
 Charlier CJ, see Joris JL  
 Chelliah RY, see Chow MYH  
 Chen EH, see Stevenson GW  
 Cheng J-K, Pan M-H, Wu K-H, Mok MS, Wei T-T. Epidural phenylephrine attenuates hypotension induced by alkalized lidocaine epidural anesthesia, 1322  
 Chestnut DH, see Bimbach DJ  
 Cheung AT, Bavaria JE, Pochettino A, Weiss SJ, Barclay DK, Stecker MM. Oxygen delivery during retrograde cerebral perfusion in humans, 8  
 Chhabra B, see Gupta R  
 Chhibber AK, see Lustik SJ  
 Chiu JW, see Sia ATH  
 Chong BK, see Chow MYH  
 Chong JL, see Sia ATH  
 Choudhry DK, see Shevchenko Y  
 Chow MYH, Liam BL, Lew TWK, Chelliah RY, Ong BC. Selecting double-lumen tubes for small patients (letter, reply), 466

- Liam BL, Thng CH, Chong BK.  
Predicting the size of a double-lumen endobronchial tube using computed tomographic scan measurements of the left main bronchus diameter, 302  
see Ti LK
- Chuang AZ, see Campos C
- Chung CW, see Kim KS
- Chung F, see Marshall SI
- Clanachan AS, see Wachowski I
- Coalson DW, see Janiszewski DJ
- Coderre TJ, see Gilon I
- Cohen D, see Gaiser RR
- Cohen DE, see Rose JB
- Cohen SE. Statistical analysis of total labor pain using the visual analog scale and application to studies of analgesic effectiveness during childbirth (letter), 1193
- Cole DJ, see Soonthon-Brant V
- Connelly NR, see Reuben SS
- Contro R, see Runza M
- Conzen PF, see Szekely A
- Cook CD, see Morgan D
- Cook DJ. Changing temperature management for cardiopulmonary bypass, 1254  
see Plöchl W
- Cooper JD, see Hogue CW Jr
- Coriat P  
see Eyraud D  
see Vivien B
- Cork RC, see Palmer CM
- Cornish PB, Greenfield LJ, O'Reilly M, Allan L. Indirect versus direct measurement of brachial plexus depth, 1113
- Cosgrove DM, see Savage RM
- Coté CJ  
see Rothschilder JL  
see Stevenson GW
- Cowan A, see Radzievsky AA
- Craig WF, see Ghoname EA
- Cramond T, see Smith MT
- Crombleholme T, see Gaiser RR
- Cruz A, see Garutti I
- Cunningham AJ, see Kelly DJ
- Custers WT, see Medici G
- Cuy R, see Rose JB
- Dahaba AA, see Oczenski W
- Dahan A, see Stienstra R
- Dahlgren G, see Blomquist H
- Dain SL, see Webster AC
- Dalman H, see Norris MC
- Daly RC, see Plöchl W
- Dang CP. Midfemoral block: a new lateral approach to the sciatic nerve (letter), 1426
- D'Angelo R, Evans E, Dean LA, Gaver R, Eisenach JC. Spinal clonidine prolongs labor analgesia from spinal sufentanil and bupivacaine, 573
- Dantzer D, see Webster AC
- Datta AK, see Gupta AK
- Datta S  
see Martin R  
see Tsen LC
- Davis PJ, Greenberg JA, Gendelman M, Fertil K. Recovery characteristics of sevoflurane and halothane in preschool-aged children undergoing bilateral myringotomy and pressure equalization tube insertion, 34
- Davis S, see Walz R
- de Boer A, see van Dongen EP
- De Cillis P, see Varrassi G
- de Menezes Lyra R. Glottis simulator (letter), 1422
- De Nicola A, see Varrassi G
- de Nucci G, see Tanus-Santos JE
- de Waal EE. The hemodynamic effects of pneumoperitoneum during laparoscopic surgery in healthy infants: assessment by continuous esophageal aortic blood flow echo Doppler (letter), 468
- Deacock SJ, see Toh KW
- Deal E, see Goldberg ME
- Dean LA, see D'Angelo R
- Denson DD, see Hord AH
- DePerio M, see Shore-Lesserson L
- D'Ercole FJ, Scott D, Bell E, Klein SM, Greengrass RA. Paravertebral blockade for modified radical mastectomy in a pregnant patient, 1351
- DeRiso BM. Management of the Operating Room. Vol. 36, No. 1 of International Anesthesiology Clinics (book review), 699
- Dexter F. Statistical analysis of total labor pain using the visual analog scale and application to studies of analgesic effectiveness during childbirth (letter, reply), 1194
- Macario A. Decrease in case duration required to complete an additional case during regularly scheduled hours in an operating room suite: a computer simulation study, 72
- Macario A, Manberg PJ, Lubarsky DA. Computer simulation to determine how rapid anesthetic recovery protocols to decrease the time for emergence or increase the phase I postanesthesia care unit bypass rate affect staffing of an ambulatory surgery center, 1053
- Visual analog scale scores for labor pain (letter, reply), 1421  
see Miura Y
- Diaz-Ruano S, see Ferrando A
- Dilrosun-Alhadi BZR, see Stienstra R
- Dirksen R, see Slappendel R
- Doğan IV, Ovali E, Eti Z, Yayci A, Göğüş FY. The *in vitro* effects of isoflurane, sevoflurane, and propofol on platelet aggregation, 432
- Domino KB, Anderson EA, Polissar NL, Posner KL. Comparative efficacy and safety of ondansetron, droperidol, and metoclopramide for preventing postoperative nausea and vomiting: a meta-analysis, 1370
- Done ML, see Lee A
- Douglas A, see Goodman EJ
- Drexler C, see Schmeling WT
- Drummond JC, see Soonthon-Brant V
- Dsida RM, see Rothschilder JL
- Dubey PK, see Sinha PK
- Dubois MY. Pain Management. Vol 6 in Atlas of Anesthesia (book review), 1430
- Dubreuil M, see Grenier B
- Dudley D, see Viscomi CM
- Dunbar PJ, Visco E, Lam AM. Craniotomy procedures are associated with less analgesic requirements than other surgical procedures, 335
- Durieux ME, see Sullivan LM
- Ebel C, see Holst D
- Ebert TJ. Is there a responsibility to disclose data used as the basis for a publication? (letter, reply), 694
- Eger EI II. Is there a responsibility to disclose data used as the basis for a publication? (letter), 693
- Halsey MJ, Harris RA, Koblin DD, Pohorille A, Sewell JC, Sonner JM, Trudell JR. Hypothesis: volatile anesthetics produce immobility by acting on two sites approximately five carbon atoms apart, 1395
- Ionescu P, Laster MJ, Gong D, Hudlicky T, Kendig JJ, Harris RA, Trudell JR, Pohorille A. Minimum alveolar anesthetic concentration of fluorinated alkanols in rats: relevance to theories of narcosis, 867
- Koblin DD, Sonner J, Gong D, Laster MJ, Ionescu P, Halsey MJ, Hudlicky T. Nonimmobilizers and transitional compounds may produce convulsions by two mechanisms, 884  
see Koblin DD

- see Ueno S
- Eichelberger JP, see Lustik SJ
- Eilers H, Kindler CH, Bickler PE. Different effects of volatile anesthetics and polyhalogenated alkanes on depolarization-evoked glutamate release in rat cortical brain slices, 1168
- Eisenach J  
see Gerancher JC  
see D'Angelo R
- El Becheiry H, see Manninen PH
- El-Schahawi-Kienzl I, see Oczenski W
- Elsharkawi NG. Simple pediatric analog sedation score (PASS) (letter), 227
- Elwell CE, see Lovell AT
- Enneking FK, see Botero M
- Erdemli E, see Kurtipek Ö
- Ergin MA, see Shore-Lesserson L
- Eriksson LJ, see Österlund Å
- Erren M  
see Booke M  
see Loick HM
- Eslambouli NR, see Celebioglu B
- Etches R, see Scott DA
- Eti Z, see Doğan IV
- Evans E, see D'Angelo R
- Eyraud D, Brabant S, Nathalie D, Fléron M-H, Gilles G, Bertrand M, Coriat P. Treatment of intraoperative refractory hypotension with terlipressin in patients chronically treated with an antagonist of the renin-angiotensin system, 980
- Ezri T, Szmuk P, Geva D. Modified esophageal detector device (letter), 691
- Fábregas N, see Suarez S
- Fagan Shier ME. A simple technique for oral fiberoptic bronchoscopy: "no more needles, doc" (letter), 695
- Fahy BG, Hasnain JU, Flowers JL, Plotkin JS, Odonkor P, Ferguson MK. Transesophageal echocardiographic detection of gas embolism and cardiac valvular dysfunction during laparoscopic nephrectomy, 500
- Fanciullo GJ, Rose RJ, Lunt PG, Whalen PK, Ross E. The state of implantable pain therapies in the United States: a nationwide survey of academic teaching programs, 1311
- Fanelli G, Casati A, Garancini P, Torri G. Nerve stimulator and multiple injection technique for upper and lower limb blockade: failure rate, patient acceptance, and neurologic complications, 847  
see Casati A
- Farber NE, Samsó E, Staunton M, Schwabe D, Schmeling WT. Dexmedetomidine modulates cardiovascular responses to stimulation of central nervous system pressor sites, 617  
see Schmeling WT
- Fassoulaki A, Sarantopoulos C, Zotou M, Karabinis G. Assessment of the level of sensory block after subarachnoid anesthesia using a pressure palpator, 398
- Faust RJ, see Kuhnert SM
- Fawcett WJ, see Toh KW
- Ferguson MK, see Fahy BG
- Ferrando A, Garutti I, Pulido C, Diaz-Ruano S, Garcia De Lucas E. Carbon dioxide embolism in an infant (letter), 227
- Ferrenbach D, see Norris MC
- Fertal K, see Davis PJ
- Fiamengo SA, see Remsen LG
- Fish DJ. The Cancer Patient. Vol. 16, No. 3 of Anesthesia Clinics of North America (book review), 699
- Fitzgerald RD, see Oczenski W
- Fitzmaurice BG, see Brodsky JB
- Fleischmann E, Akça O, Wallner T, Arkiliç CF, Kurz A, Hickel RS, Zimpfer M, Sessler DI. Onset time, recovery duration, and drug cost with four different methods of inducing general anesthesia, 930
- Fleisher LA  
see Lehmann HP  
see Nelson JB
- Fléron M-H, see Eyraud D
- Fletcher H, see Myles PS
- Flick R, Martin LD. Twelfth annual meeting of the Society for Pediatric Anesthesia, Orlando, Florida, October 16, 1998, 955
- Flowers JL, see Fahy BG
- Floyd H, see Gerancher JC
- Fogel ST, Leighton B. Epidural analgesia and cesarean delivery: what is the relationship? (letter, reply), 964  
see Norris MC
- Fontana JL, see Musser JB
- Forrest AP, see Found P
- Fortney JT, Gan TJ, Glass PSA. Ondansetron versus droperidol for postoperative nausea and vomiting (letter, reply), 465
- Fosse E, see Klaastad Ø
- Foster E, see Lambert A-S
- Found P, Forrest AT. Vaporizer level obstruction detected by anesthetic vapor analysis (letter), 469
- Fradis M, see Gaitini LA
- Francis S, see Shore-Lesserson L
- Frank SM, Nguyen JM, Garcia CM, Barnes RA. Temperature monitoring practices during regional anesthesia, 373  
see Nelson JB
- Frass M, see Knapp S
- Fredman B, Nun MB, Zohar E, Iraqi G, Shapiro M, Gepstein R, Jedeikin R. Epidural steroids for treating "failed back surgery syndrome": is fluoroscopy really necessary?, 367
- Zohar E, Golan E, Tillingier M, Bernheim J, Jedeikin R. Diclofenac does not decrease renal blood flow or glomerular filtration in elderly patients undergoing orthopedic surgery, 149
- Freedman JM, Rudow MP. Bilateral buttock and leg pain after lidocaine epidural anesthesia (letter), 1188
- Freid EB, see Valley RD
- Frenette L. Transfusion requirements in orthotopic liver transplantation (letter, reply), 692
- Frink BA, see Lehmann HP
- Fujii Y, Saitoh Y, Tanaka H, Toyooka H. Granisetron/dexamethasone combination for reducing nausea and vomiting during and after spinal anesthesia for cesarean section, 1346
- Saitoh Y, Tanaka H, Toyooka H. Preoperative oral antiemetics for reducing postoperative vomiting after tonsillectomy in children: granisetron versus perphenazine, 1298
- Fujimoto S, see Sumida M
- Fujioka S, Ono Y, Kangawa K, Okada K. Plasma concentration of adrenomedullin is increased in hemorrhagic shock in dogs, 326
- Fujita H, see Soga K
- Fukumoto M, see Adachi T
- Furuya H  
see Inoue S  
see Kawaguchi M
- Furuya R, Oka K, Watanabe I, Kamiya Y, Itoh H, Andoh T. The effects of ketamine and propofol on neuronal nicotinic acetylcholine receptors and P<sub>2u</sub> purinoceptors in PC12 cells, 174  
see Watanabe I
- Gabriele S, see Reich DL
- Gaiser RR, Kurth CD, Cohen D, Crombleholme T. The cesarean delivery of a twin gestation under



- 2 minimum alveolar anesthetic concentration isoflurane: one normal and one with a large neck mass, 584
- Gaitini LA, Vaida SJ, Somri M, Fradis M, Ben-David B. Fiberoptic-guided airway exchange of the esophageal-tracheal Combitube in spontaneously breathing versus mechanically ventilated patients, 193
- Gajraj NM, see Ghoname EA
- Galbraith JC, see Wachowski I
- Galea M, see Beilin Y
- Galinkin JL, see Janiszewski DJ
- Gan TJ, Bennett-Guerrero E, Phillips-Bute B, Wekeling H, Moskowitz DM, Olufolabi Y, Konstadt SN, Bradford C, Glass PSA, Machin SJ, Mythen MG. Hextend, a physiologically balanced plasma expander for large volume use in major surgery: a randomized phase III clinical trial, 992
- Lubarsky DA. Cost effectiveness ratio: an often misunderstood term (letter), 1191
- Ganapathy S, see Mulroy MF
- Ganjoo P, see Schmeling WT
- Garancini P, see Fanelli G
- Garcia CM, see Frank SM
- Garcia de Lucas E  
see Ferrando A  
see Garutti I
- Garry B. Rediscovering the obvious (letter, reply), 1427
- Garutti I, Quintana B, Olmedilla L, Cruz A, Barranco M, Garcia de Lucas E. Arterial oxygenation during one-lung ventilation: combined versus general anesthesia, 494  
see Ferrando A
- Gat TJ, see Fortney JT
- Gaver R, see D'Angelo R
- Geisslinger G, see Koppert W
- Gendelman M, see Davis PJ
- Gentili M, Bernard J-M, Bonnet F. Adding clonidine to lidocaine for intravenous regional anesthesia prevents tourniquet pain, 1327
- Georges LS, see Valley RD
- Gepstein R, see Fredman B
- Gerancher JC, Floyd H, Eisenach J.  
Determination of an effective dose of intrathecal morphine for pain relief after cesarean delivery, 346  
Valuing the work performed by anesthesiology residents (letter), 964  
see Habibi A
- Gerber H, see Konrad C
- Geva D, see Ezri T
- Ghoname EA, Craig WF, White PF, Ahmed HE, Hamza MA, Gajraj NM, Vakharia AS, Noe CE. The effect of stimulus frequency on the analgesic response to percutaneous electrical nerve stimulation in patients with chronic low back pain, 841
- Gibbs NM, see Weightman WM
- Gibson CS, see Reuben SS
- Gielen MJM, see Slappendel R
- Gilbert TB, Goodsell CW, Krasna MJ. Bronchial rupture by a double-lumen endobronchial tube during staging thoracoscopy, 1252
- Giller G, see Eyraud D
- Gilron I, Quirion R, Coderre TJ. Pre-versus postinjury effects of intravenous GABAergic anesthetics on formalin-induced fos immunoreactivity in the rat spinal cord, 414
- Girling KJ, Bedford NM, Spendlove JL, Mahajan RP. Assessing neuromuscular block at the larynx: the effect of change in resting cuff pressure and a comparison with video imaging in anesthetized humans, 426
- Giunta F, see Varrassi G
- Glass PSA  
see Fortney JT  
see Gan TJ
- Gluncic I, see Kapural L
- Golan E, see Fredman B
- Goldberg ME, Cantillo J, Gratz I, Deal E, Vekeman D, McDougall R, Afshar M, Zafeiridis A, Larijani G. Dose of compound A, not sevoflurane, determines changes in the biochemical markers of renal injury in healthy volunteers, 437
- Goldstone JC, see Lovell AT
- Gollmann G, see Kapral S
- Gong D  
see Eger EI II  
see Koblin DD
- Goodman EJ, Hudson IM, Douglas A. Desflurane vaporizer uses minimal electricity (letter), 1189
- Goodsell CW, see Gilbert TB
- Gordon W, see Reich DL
- Gossler K, see Palmer CM
- Gothard JWW. Endobronchial intubation in infants and small children (letter), 229
- Goto T, see Ichinose F
- Gotta AW. The Laryngeal Mask Airway. Vol. 36, No. 2 of International Anesthesiology Clinics (book review), 697
- Gottschall R, see Klein U
- Goudsouzian NG, see Nishanian EV
- Gourdange PA, see Broka SM
- Gögüş FY, see Doğan IV
- Grabenwöger M, see Rajek A
- Grady RE, Horlocker TT, Brown RD, Maxson PM, Schroeder DR. Neurologic complications after placement of cerebrospinal fluid drainage catheters and needles in anesthetized patients: implications for regional anesthesia, 388
- Gratz I, see Goldberg ME
- Greacen M, see Wachowski I
- Green NM. Gwenifer Wilson, M.D., 1916-1998: Author, Historian, Laureate (obituary), 690
- Greenberg JA, see Davis PJ
- Greenberg RS, Prichard CB, Jacquin KJ. Repeated use of the cuffed oropharyngeal airway in an infant for radiation therapy (letter), 1421
- Greenfield LJ, see Cornish PB
- Greengrass RA  
see D'Ercole FJ  
see Mulroy MF
- Greilich PE, see Whitten CW
- Greim C-A, Brederlau J, Kraus I, Apfel C, Thiel H, Roewer N. Transnasal transesophageal echocardiography: a modified application mode for cardiac examination in ventilated patients, 306
- Grenier B, Verchère E, Mesli A, Dubreuil M, Siao D, Vandendriessche M, Calès J, Maurette P. Capnography monitoring during neurosurgery: reliability in relation to various intraoperative positions, 43
- Gries A, Bode C, Gross S, Peter K, Böhrer H, Martin E. The effect of intravenously administered magnesium on platelet function in patients after cardiac surgery, 1213
- Grocott HP, Sheng H, Miura Y, Sarraf-Yazdi S, Mackensen GB, Pearlstein RD, Warner DS. The effects of aprotinin on outcome from cerebral ischemia in the rat, 1
- Gross JB. Total Intravenous Anesthesia (book review), 698
- Gross S, see Gries A
- Gruber EM  
see Tschernko EM  
see Rajek A

- Gueugniaud P-Y, Pannetier J-C. The hemodynamic effects of pneumoperitoneum during laparoscopic surgery in healthy infants: assessment by continuous esophageal aortic blood flow echo Doppler (letter, reply), 469
- Gugel M, see Klein U
- Gupta AK, Hutchinson PJ, Al-Rawi P, Gupta S, Swart M, Kirkpatrick PJ, Menon DK, Datta AK. Measuring brain tissue oxygenation compared with jugular venous oxygen saturation for monitoring cerebral oxygenation after traumatic brain injury, 549  
see Summors AC
- Gupta R, Singhal SK, Rattan KN, Chhabra B. Endobronchial intubation in infants and small children (letter, reply), 229
- Gupta S, see Gupta AK
- Gustafsson U, see Scott DA
- Gwartz KH, Young JV, Byers RS, Alley C, Levin K, Walker SG, Stoelting RK. The safety and efficacy of intrathecal opioid analgesia for acute postoperative pain: seven years' experience with 5969 surgical patients at Indiana University Hospital, 599
- Habibi A, Schmeising C, Gerancher JC. Interscalene regional anesthesia in the prevention of autonomic hyperreflexia in a quadriplegic patient undergoing shoulder surgery, 98
- Hagelberg A, see Lauwers M
- Hagemann O, see Booke M
- Haider W, see Tschernko EM
- Hakim TS, see Thomas PS
- Hall DB. Takes a lickin' and keeps on tickin' (letter), 1424
- Hall SC, see Stevenson GW
- Hall SR, Wang L, Milne B, Loomis C. Activation of the Rostral ventrolateral medulla in an acute anesthetized rodent strychnine model of allodynia, 1125
- Halliwell R, see Scott DA
- Halpern P, see Hochhauser E
- Halsey MJ  
see Eger EI II  
see Koblin DD
- Hamoir EE, see Joris JL
- Hamu Y, Kanmura Y, Tsuneyoshi I, Yoshimura N. The effects of vasopressin on endotoxin-induced attenuation of contractile responses in human gastroepiploic arteries *in vitro*, 542
- Hamza MA, see Ghoname EA
- Hansen D, Hansen S. The effects of three graded doses of meperidine for spinal anesthesia in African men, 827
- Syben R, Vargas O, Spies C, Welte M. The alveolar-arterial difference in oxygen tension increases with temperature-corrected determination during moderate hypothermia, 538
- Hansen S, see Hansen D
- Hargett MJ, see Kahn RL
- Harris RA  
see Eger EI II  
see Ueno S
- Hartstein GM, see Joris JL
- Harvey SC, see Overdyk FJ
- Hase K, Meguro K, Nakamura T. Assessment of renal effects of sevoflurane in elderly patients using urinary markers (letter), 1426
- Hasnain JU, see Fahy BG
- Hass D, see Lien CA
- Hatori F, see Ichinose F
- Hayashi M, see Adachi T
- Hayashi Y, see Inoue S
- Haynes GR, see Burt N
- Heavner JE, Rosenberg P. Propofol for lidocaine-induced seizures (letter), 1193
- Shi B, Pitkänen M. Nitric oxide synthesis inhibition modifies the cardiotoxicity of tetracaine and lidocaine, 717  
see Nyström EUM
- Hedner T, see Valsson F
- Heierson A, see Mulroy MF
- Heindl B, see Szekely A
- Hendolin H, see Kokki H
- Hibbard M, see Reich DL
- Hickle RS, see Fleischmann E
- Hildebrand KL, see Williams EL
- Hildebrandt U, see Vollmar B
- Hirakawa M, see Otsuka F
- Hirose Y, see Kawahito S
- Hirota K, see Wakasugi M
- Ho S-T  
see Wang J-J  
see Wu C-T
- Hobbahn J, see Keyl C
- Hobbs GJ, see Tighe KE
- Hochhauser E, Halpern P, Zolotarsky V, Krasnov T, Sulkes J, Vidne B. Isoflurane and sodium nitroprusside reduce the depressant effects of protamine sulfate on isolated ischemic rat hearts, 710
- Hochmuth K, see Keyl C
- Hodgson PS, Neal JM, Pollock JE, Liu SS. The neurotoxicity of drugs given intrathecally (spinal), 797
- Hoerauf KH, Wallner T, Akça O, Taslimi R, Sessler DI. Exposure to sevoflurane and nitrous oxide during four different methods of anesthetic induction, 925
- Hofbauer R, see Knapp S
- Hofstadter MB, see Kain ZN
- Hogue CW Jr, Stamos T, Winters KJ, Moulton M, Krucylak PE, Cooper JD. Acute myocardial infarction during lung volume reduction surgery, 332
- Holst D, Möllmann M, Schymroszczyk B, Ebel C, Wendt M. No risk of metal toxicity in combined spinal-epidural anesthesia, 393
- Hönemann CW, see Sullivan LM
- Hooda S. Where wrong practice is a necessity (letter), 233
- Hop WCJ, see Medici G
- Hoppe W, see Norris MC
- Hoppert S, see Puura AIE
- Hord AH, Denson DD, Huerkamp MJ, Seiler JG III. Changes in rat paw perfusion after experimental mononeuropathy: assessment by laser Doppler fluxmetry, 103
- Horlocker TT, Kufner RP, Bishop AT, Maxson PM, Schroeder DR. The risk of persistent paresthesia is not increased with repeated axillary block, 382  
see Grady RE  
see Wedel DJ
- Horn A, see Lauwers M
- Horn B, see Stevenson GW
- Hornstein MD, see Martin R
- Hosain SI, see Thomas PS
- Houltz E, see Valsson F
- Hrazdil J, see Wachowski I
- Hubert BM, see Joris JL
- Hudlicky T  
see Eger EI II  
see Koblin DD
- Hudson IM, see Goodman EJ
- Huerkamp MJ, see Hord AH
- Huffnagle S, Norris MC. The influence of epidural needle bevel on spread of sensory blockade in the laboring parturient (letter, reply), 962
- Huizar K, see Scott DA
- Hunt JO, see Myles PS
- Huntington JH, Malviya S, Voepel-Lewis T, Lloyd TR, Massey KD. The effect

- of a right-to-left intracardiac shunt on the rate of rise of arterial and end-tidal halothane in children, 759
- Hutchinson PJ, see Gupta AK
- Iatrou CC, see Leslie K
- Ichinose F, Uezono S, Muto R, Uchida H, Hatori F, Terui K, Niimi Y, Goto T, Nakata Y, Morita S. Platelet hyporeactivity in young infants during cardiopulmonary bypass, 258
- Ikeda K, see Ikeda T
- Ikeda T, Sessler DI, Kikura M, Kazama T, Ikeda K, Sato S. Less core hypothermia when anesthesia is induced with inhaled sevoflurane than with intravenous propofol, 921
- Ikeuchi Y, see Kawahito S
- Inaba H, see Sumida M
- Inagaki Y, see Sá Rêgo MM
- Inoue S, Hayashi Y, Ohnishi Y, Kikumoto K, Minamino N, Kangawa K, Matsuo H, Furuya H, Kuro M. Cerebral production of adrenomedullin after hypothermic cardiopulmonary bypass in adult cardiac surgical patients, 1030
- see Kawaguchi M
- Ionescu P
- see Eger EI II
- see Koblin DD
- Iraqi G, see Fredman B
- Isawa E, see Sumida M
- Ischia S, see Varrassi C
- Ishikawa S. Is continuous intraarterial blood gas monitoring reliable during one-lung ventilation? (letter), 1192
- Islander G, Twetman ER. Comparison between the European and North American protocols for diagnosis of malignant hyperthermia susceptibility in humans, 1155
- Ito Y, see Wakasugi M
- Itoh H
- see Furuya R
- see Watanabe I
- Ivy R, see Whitten CW
- Izenberg N
- see Cassidy JF Jr
- see Miller KM
- Jacquin KJ, see Greenberg RS
- James RL, see Politis GD
- Jandrasits O
- see Kapral S
- see Tschernko EM
- Janiszewski DJ, Galinkin JL, Klock PA, Coalson DW, Pardo H, Zacny JP. The effects of subanesthetic concentrations of sevoflurane and nitrous oxide, alone and in combination, on analgesia, mood, and psychomotor performance in healthy volunteers, 1149
- Jantsch U, see Rajek A
- Jantsch-Watzinger U, see Tschernko EM
- Janzen V, see Webster AC
- Jedeikin R, see Fredman B
- Jellinek H, see Oczenski W
- Jellish WS, Blakeman B, Warf P, Slogoff S. Somatosensory evoked potential monitoring used to compare the effect of three asymmetric sternal retractors on brachial plexus function, 292
- Jocson JC, see Shevchenko Y
- Johmura S, see Asai T
- Jöhr M, Sossai R. Colonic puncture during ilioinguinal nerve block in a child, 1051
- Jolly DT, see Wachowski I
- Jones K, see Leslie K
- Jönsson A, see Mattsson U
- Jontell M, see Mattsson U
- Joo HS, Rose DK. The intubating laryngeal mask airway with and without fiberoptic guidance, 662
- Joris JL, Hamoir EE, Hartstein GM, Meurisse MR, Hubert BM, Charlier CJ, Lamy ML. Hemodynamic changes and catecholamine release during laparoscopic adrenalectomy for pheochromocytoma, 16
- Joucken KL, see Broka SM
- Jun JH, see Kim KS
- Junker R, see Loick HM
- Kadoya T
- see Nijijima K
- see Seto A
- Kahn RL, Hargett MJ.  $\beta$ -Adrenergic blockers and vasovagal episodes during shoulder surgery in the sitting position under interscalene block, 378
- Kain ZN. Perioperative information and parental anxiety: the next generation (editorial), 237
- Wang SM, Mayes LC, Caramico LA, Hofstadter MB. Distress during the induction of anesthesia and postoperative behavioral outcomes, 1042
- Kal JE, Vergroesen I, van Wezel HB. The effect of nitroglycerin on pacing-induced changes in myocardial oxygen consumption and metabolic coronary vasodilation in patients with coronary artery disease, 271
- Kalkman CJ
- see Kochs E
- see Ubags LH
- Kamiya Y
- see Furuya R
- see Watanabe I
- Kanbak M. Massive intraoperative pulmonary embolism: with succinylcholine or not? (letter), 1423
- Kanda T, Ohta Y, Kida A, Kemmotsu O. The effect of  $\alpha_2$ -adrenergic drugs on the activity of neurons in the rat nucleus raphe magnus *in vitro*, 459
- Kangawa K
- see Fujioka S
- see Inoue S
- Kanmura Y, see Hamu Y
- Kapral S, Gollmann G, Bachmann D, Prohaska B, Likar R, Jandrasits O, Weinstabl C, Lehofer F. The effects of thoracic epidural anesthesia on intraoperative visceral perfusion and metabolism, 402
- Gollmann G, Walt B, Likar R, Sladen RN, Weinstabl C, Lehofer F. Tramadol added to mepivacaine prolongs the duration of an axillary brachial plexus blockade, 853
- Kapural L, Sprung J, Gluncic I, Kapural M, Andelinovic S, Primorac D, Schoenwald PK. Tracheo-innominate artery fistula after tracheostomy, 777
- Kapural M, see Kapural L
- Karabinis G, see Fassoulaki A
- Karamichali EA, see Mentzelopoulos SD
- Karzai W, see Klein U
- Kastner J, see Rajek A
- Kawaguchi M, Inoue S, Sakamoto T, Kawaraguchi Y, Furuya H, Sakaki T. The effects of prostaglandin  $E_1$  on intraoperative temperature changes and the incidence of postoperative shivering during deliberate mild hypothermia for neurosurgical procedures, 446
- Kawahito S, Kitahata H, Tanaka K, Ikeuchi Y, Kimura H, Hirose Y, Oshita S. Intraoperative management of a pediatric patient undergoing cardiac tumor resection with the aid of transesophageal and epicardial echocardiography, 1048
- Kawamata M, see Sumida M
- Kawaraguchi Y, see Kawaguchi M
- Kayne RD, see Reich DL
- Kazama T, see Ikeda T
- Keller C, Brimacombe J. Mucosal pressure, mechanism of seal, airway sealing pressure, and anatomic position for the disposable versus reusable

- laryngeal mask airways, 1418
- Brimacombe J. The influence of head and neck position on oropharyngeal leak pressure and cuff position with the flexible and the standard laryngeal mask airway, 913
- Brimacombe J, Rädler C, Pühringer F. Do laryngeal mask airway devices attenuate liquid flow between the esophagus and pharynx? A randomized, controlled cadaver study, 904
- Kelly DJ, Walsh F, Norman GS, Cunningham AJ. The effects of midazolam on pure tone audiometry, speech audiometry, and audiological reaction times in human volunteers, 1064
- Kelly PJ, see Soonthon-Brant V
- Kemmotsu O, see Kanda T
- Kendig JJ, see Eger EI II
- Keyl C, Lemberger P, Palitzsch K-D, Hochmuth K, Liebold A, Hobbhahn J. Cardiovascular autonomic dysfunction and hemodynamic response to anesthetic induction in patients with coronary artery disease and diabetes mellitus, 985
- Khalil SN, see Campos C
- Khorasani A, Candido KD, Saatee S, Khorasani A. The relationship between dynamic compliance and inspiratory flow (letter), 465
- see Khorasani A
- Kida A, see Kanda T
- Kikumoto K, see Inoue S
- Kikura M, see Ikeda T
- Kim KS, Shim JC, Jun JH, Lee KH, Chung CW. Rabbits treated with chronic isepamicin are resistant to mivacurium and rocuronium, 654
- Kimura H, see Kawahito S
- Kindler CH, see Eilers H
- Kirkegard-Nielsen H, see Talke PO
- Kirkpatrick PJ, see Gupta AK
- Kirnö K, see Valsö F
- Kirvelä M, see Laviola S
- Kissin I, see Vinik HR
- Kitahata H, see Kawahito S
- Klaastad Ø, Lilleås FG, Røtnes JS, Breivik H, Fosse E. Magnetic resonance imaging demonstrates lack of precision in needle placement by the infraclavicular brachial plexus block described by Raj et al., 593
- Klein SM, see D'Ercole FJ
- Klein U, Karzai W, Gottschall R, Gugel M, Bartel M. Respiratory gas monitoring during high-frequency jet ventilation for tracheal resection using a double-lumen jet catheter, 224
- Klepcko W, see Tschernko EM
- Klock PA, see Janiszewski DJ
- Knapp S, Kofler J, Stoiser B, Thalhammer F, Burgmann H, Posch M, Hofbauer R, Stanzel M, Frass M. The assessment of four different methods to verify tracheal tube placement in the critical care setting, 766
- Kobayashi T, see Ohmura S
- Koblin DD, Laster MJ, Ionescu P, Gong D, Eger EI II, Halsey MJ, Hudlicky T. Polyhalogenated methyl ethyl esters: solubilities and anesthetic properties, 1161
- see Eger EI II
- Kochs E, Kalkman CJ, Thornton C, Newton D, Bischoff P, Kuppe H, Abke J, Konecny E, Nahm W, Stockmanns G. Middle latency auditory evoked responses and electroencephalographic derived variables do not predict movement to noxious stimulation during 1 minimum alveolar anesthetic concentration isoflurane/nitrous oxide anesthesia, 1412
- see Schneck H
- Koelman JH, see Ubags LH
- Koetter KP, see Boldt J
- Kofler J, see Knapp S
- Koizumi K, see Shiga T
- Kokki H, Tuovinen K, Hendolin H. The effect of intravenous ketoprofen on postoperative epidural sufentanil analgesia in children, 1036
- Konecny E, see Kochs E
- Konrad C, Markl T, Schuepfer G, Gerber H, Tschopp M. The effects of *in vitro* hemodilution with gelatin, hydroxyethyl starch, and lactated Ringer's solution on markers of coagulation: an analysis using SONOCLOT, 483
- Konstadt SN, see Gan TJ
- Kopp VJ, see Valley RD
- Koppert W, Likar R, Geisslinger G, Zeck S, Schmelz M, Sittl R. Peripheral antihyperalgesic effect of morphine to heat, but not mechanical, stimulation in healthy volunteers after ultraviolet-B irradiation, 117
- Korttilä KT, see Vakkuri AP
- Kosaka Y, see Sakura S
- Kozek-Langenecker SA, Semsroth M. Effective dose of isoproterenol-containing test dose in anesthetized children (letter, reply), 1189
- Krasna MJ, see Gilbert TB
- Krasnov T, see Hochhauser E
- Kraus I, see Greim C-A
- Krenn H, see Oczenski W
- Krensavage TJ. Saline solution as lubrication to manipulate a stuck fiberoptic bronchoscope (letter), 965
- Kritzinger M, see Tschernko EM
- Krucylak PE, see Hogue CW Jr
- Kufner RP, see Horlocker TT
- Kuhnert SM, Faust RJ, Berge KH, Piepgras DG. Postoperative macroglossia: report of a case with rapid resolution after extubation of the trachea, 220
- Kumle B, see Suttner S
- Kuppe H, see Kochs E
- Kuro M, see Inoue S
- Kurth CD, see Gaiser RR
- Kurtipek Ö, Oral M, Başar HT, Aşık İ, Ateş Y, Kuzu I, Erdemli E, Ökten F, Tüzünler F. Histopathologic changes after repetitive peridural administration of metoclopramide in dogs, 100
- Kurz A, see Fleischmann E
- Kushizaki H, see Sakura S
- Kuzu I, see Kurtipek Ö
- Lakhotia R, Pandey CK, Tewari P. Arterialization of the dorsum vein on the hand (letter), 1425
- Lam AM, see Dunbar PJ
- Lam J, see Lehmann HP
- Lambert A-S, Miller JP, Merrick SH, Schiller NB, Foster E, Muhiudeen-Russell I, Cahalan MK. Improved evaluation of the location and mechanism of mitral valve regurgitation with a systematic transesophageal echocardiography examination, 1205
- see Lutz JT
- Lamy ML, see Joris JL
- Landesburg G, Zhou W, Aversano T. Tachycardia-induced subendocardial necrosis in acutely instrumented dogs with fixed coronary stenosis, 973
- Langemeijer HJ, see van Dongen EP
- Langeron O, see Vivien B
- Larijani G, see Goldberg ME
- Laster MJ
- see Eger EI II
- see Koblin DD
- Lauwers M, Camu F, Breivik H, Hagelberg A, Rosen M, Sneyd R, Horn A,

- Noronha D, Shaikh S. The safety and effectiveness of remifentanyl as an adjunct sedative for regional anesthesia, 134
- Laviola S, Kirvelä M, Spoto M-R, Tschuor S, Alon E. Pneumocephalus with intense headache and unilateral pupillary dilatation after accidental dural puncture during epidural anesthesia for cesarean section, 582
- Lawson P, see Ramanathan J
- Lee A, Done ML. The use of nonpharmacologic techniques to prevent postoperative nausea and vomiting: a meta-analysis, 1362
- Lee C, see Sun X-g
- Lee KH, see Kim KS
- Lee M, see Macario A
- Lee S-C, see Wang J-J
- Lee TL, see Ti LK
- Lehmann HP, Fleisher LA, Lam J, Frink BA, Bass EB. Patient preferences for early discharge after laparoscopic cholecystectomy, 1280
- Lehofer F, see Kapral S
- Leighton BL, see Fogel ST
- Lemak NA, see Campos C
- Lemberger P, see Keyl C
- Lenhardt R, see Rajek A
- Lepri A, see Sia S
- Lertakyananee J, see Visalyaputra S
- Leslie K, Iatrou CC, Jones K, Beemer GH. Common peroneal nerve stimulation for neuromuscular monitoring: evaluation in awake volunteers and anesthetized patients, 197
- Levin K, see Gwartz KH
- Lew TWK, see Chow MYH
- Li BW, see Xue FS
- Li C-Y, see Wu C-T
- Liam BL, see Chow MYH
- Liao X, see Xue FS
- Licker M, Schweizer A. Vasopressin and postcardiopulmonary bypass refractory hypotension (letter), 695
- Liebold A, see Keyl C
- Lien CA, Belmont MR, Abalos A, Hass D, Savarese JJ. The nature of spontaneous recovery from mivacurium-induced neuromuscular block, 648
- Likar R  
see Kapral S  
see Koppert W
- Lilleås FG, see Klaastad Ø
- Lindahl SGE  
see Österlund Å  
see Sjögren D
- Lindgren L, see Vakkuri AP
- Lindsay DR. False comfort from a pulse oximeter (letter), 1428
- Link J. Principles of Anesthetic Techniques and Anesthetic Emergencies. Atlas of Anesthesia Series. Vol. 4 (book review), 968
- Liu JH, see Xue FS
- Liu PL. Simulators in Anesthesiology Education (book review), 472
- Liu SS, see Hodgson PS
- Liu S-T, see Wu C-T
- Liu Y-C, see Wang J-J
- Lloyd TR, see Huntington JH
- Locke MM, Spiekermann BF, Rich GF. Trigemino-vagal reflex during repair of a nasal fracture under general anesthesia, 1183
- Loick HM, Schmidt C, Van Aken H, Junker R, Erren M, Berendes E, Rolf N, Meißner A, Schmid C, Scheld HH, Möllhoff T. High thoracic epidural anesthesia, but not clonidine, attenuates the perioperative stress response via sympatholysis and reduces the release of troponin T in patients undergoing coronary artery bypass grafting, 701
- Loomis C, see Hall SR
- Lovell AT, Owen-Reece H, Elwell CE, Smith M, Goldstone JC. Continuous measurement of cerebral oxygenation by near infrared spectroscopy during induction of anesthesia, 554
- Lozano AM, see Manninen PH
- Lubarsky DA. Valuing the work performed by anesthesiology residents (letter, reply), 964  
see Dexter F  
see Gan TJ
- Lucchinetti E, see Zaugg M
- Lundin S, see Valsöon F
- Lunt PG, see Fanciullo GJ
- Luo LK, see Xue FS
- Lurie SD, see Reuben SS
- Lustik SJ, Eichelberger JP, Chhibber AK, Bronsther O. Perioperative torsade de pointes ventricular tachycardia (letter, reply), 966
- Lutz JT, Lambert A-S, Rouine-Rapp K. Damage to pilot cuff tubing during intraoperative transesophageal echocardiography (letter), 1187
- Lynch C III, see McDowell TS
- Macario A, Weinger M, Truong P, Lee M. Which clinical anesthesia outcomes are both common and important to avoid? The perspective of a panel of expert anesthesiologists, 1085  
see Brodsky JB  
see Dexter F  
see Torralva PR
- Machin SJ, see Gan TJ
- Maciolek H, see Reuben SS
- Maciulla JE, see Palmer CM
- Mackensen GB  
see Grocott HP  
see Miura Y
- Maekawa N, see Nishina K
- Mahajan RP. The effects of propofol on cerebral blood flow velocity and cerebral oxygen extraction during cardiopulmonary bypass (letter), 231  
see Girling KJ
- Mahon BD, see Weightman WM
- Makino H, see Otsuka F
- Maleck W, see Boldt J
- Malinovsky J-M, see Bouaziz H
- Mallios C, see Medici G
- Malviya S  
see Huntington JH  
see Tait AR
- Manberg PJ, see Dexter F
- Manecke GR Jr, Nemirov MA, Bicker AA, Adsumelli RN, Poppers PJ. The effect of halothane on the amplitude and frequency characteristics of heart sounds in children, 263
- Manninen PH, Burke SJ, Wennberg R, Lozano AM, El Beheiry H. Intraoperative localization of an epileptogenic focus with alfentanil and fentanyl, 1101  
see Puura AIE
- Manspeizer HE, see Shore-Lesserson L
- Mantero S, see Runza M
- Marcus MAE. Isoproterenol and epidural analgesia (letter, reply), 1190  
see Brodner G
- Mares P  
see Rajek A  
see Tschernko EM
- Marinangeli F, see Varrasi G
- Markl T, see Konrad C
- Marrin CAS, see Rassias AJ
- Marshall SI, Chung F. Discharge criteria and complications after ambulatory surgery, 508
- Marsland C, see Scott DA
- Martin E, see Gries A
- Martin LD, see Flick R
- Martin R, Tsen LC, Tzeng G, Hornstein MD, Datta S. Anesthesia for *in vitro* fertilization: the addition of fentanyl to 1.5% lidocaine, 523



- Martínez-Pérez A, see Candela-Toha AM  
 Massey KD, see Huntington JH  
 Matsumoto M, see Wakamatsu H  
 Matsuo H, see Inoue S  
 Matsuzaki T, see Yamakage M  
 Matta BF, see Summors AC  
 Mattsson U, Cassuto J, Jontell M, Jönsson A, Sinclair R, Tarnow P. Digital image analysis of erythema development after experimental thermal injury to human skin: effect of postburn topical local anesthetics (EMLA), 1131  
 Maurette P, see Grenier B  
 Maxson PM  
   see Grady RE  
   see Horlocker TT  
 Mayes LC, see Kain ZN  
 Mazoit JX, see Bouaziz H  
 McAdoo J, see McGinley J  
 McCarthy RJ, see Tuman KJ  
 McCormick CI, see Remsen LG  
 McCormick SA, see Williams EL  
 McDougall R, see Goldberg ME  
 McDowell TS, Pancrazio JJ, Barrett PQ, Lynch C III. Volatile anesthetic sensitivity of T-type calcium currents in various cell types, 168  
 McGinley J, McAdoo J. Airway adjunct to an unanticipated difficult airway (letter), 467  
 McRae VA, see Shevchenko Y  
 Medici G, Mallios C, Custers WT, van Meerbeek JP, Verhoeven GT, Hop WCJ. Anesthesia for endobronchial laser surgery: a modified technique, 298  
 Meguro K, see Hase K  
 Meißner A, see Loick HM  
 Mekhail NA. Management of Acute and Chronic Pain (book review), 967  
 Menger MD, see Vollmar B  
 Menon DK, see Gupta AK  
 Mentzelopoulos SD, Tsitsika MV, Karamichali EA. Difficult airway management with balloon inflation (letter), 1425  
 Merrick SH, see Lambert A-S  
 Merridew G, see Scott DA  
 Mertes N, see Brodner G  
 Mesli A, see Grenier B  
 Meurisse MR, see Joris JL  
 Meyer DJ, see Tobias JD  
 Mikawa K, see Nishina K  
 Miller JP, see Lambert A-S  
 Miller KM, Wysocki T, Cassady JF Jr, Cancell D, Izenberg N. Validation of measures of parents' preoperative anxiety and anesthesia knowledge, 251  
   see Cassady JF Jr  
 Milne B, see Hall SR  
 Minamino N, see Inoue S  
 Miura M, see Otsuka F  
 Miura Y, Mackensen GB, Nellgård B, Pearlstein RD, Bart RD, Dexter F, Warner DS. Effects of isoflurane, ketamine, and fentanyl/N<sub>2</sub>O on concentrations of brain and plasma catecholamines during near-complete cerebral ischemia in the rat, 787  
   see Grocott HP  
 Miyazaki Y, Adachi T, Utsumi J, Shichino T, Segawa H. Xenon has greater inhibitory effects on spinal dorsal horn neurons than nitrous oxide in spinal cord transected cats, 893  
 Mizutani K, see Shogase A  
 Mok MS, see Cheng J-K  
 Möllhoff T  
   see Brodner G  
   see Loick HM  
 Möllmann M, see Holst D  
 Momota Y, see Artru AA  
 Mongan PD, see Musser JB  
 Moore DC. Label and labeling are not interchangeable (letter), 467  
 Morell RC, see Politis GD  
 Moreno H Jr, Tanus-Santos E. Nitric oxide inhalation during massive pulmonary embolism (letter), 1188  
   see Tanus-Santos JE  
 Morgan D, Cook CD, Smith MA, Picker MJ. An examination of the interactions between the antinociceptive effects of morphine and various  $\mu$ -opioids: the role of intrinsic efficacy and stimulus intensity, 407  
 Mori T, see Yamakage M  
 Morimont CJ, see Broka SM  
 Morita K, see Otsuka F  
 Morita S, see Ichinose F  
 Morley-Forster PK, see Webster AC  
 Morshuis WJ, see van Dongen EP  
 Moskowitz DM, see Gan TJ  
 Moulton M, see Hogue CW Jr  
 Muhiudeen-Russell I, see Lambert A-S  
 Mulroy MF, Greengrass R, Ganapathy S, Chan V, Heierson A. Sameridine is safe and effective for spinal anesthesia: a comparative dose-ranging study with lidocaine for inguinal hernia repair, 815  
 Munro HM, see Tait AR  
 Murphy D, see Scott DA  
 Musser JB, Fontana JL, Mongan PD. The anesthetic and physiologic effects of an intravenous administration of a halothane lipid emulsion (5% vol/vol), 671  
 Muto R, see Ichinose F  
 Myles PS, Hunt JO, Nightingale CE, Fletcher H, Beh T, Tanil D, Nagy A, Rubinstein A, Ponsford JL. Development and psychometric testing of a quality of recovery score after general anesthesia and surgery in adults, 83  
 Mythen MG, see Gan TJ  
 Naganuma K, see Nishiyama M  
 Nagi H, Brown PC. Undetected hole in a laryngeal mask (letter), 232  
 Naguib SS, see Campos C  
 Nagy A, see Myles PS  
 Nahm W, see Kochs E  
 Naimi C, see Raby KE  
 Nakakimura K, see Wakamatsu H  
 Nakamura T, see Hase K  
 Nakao S, see Shinomura T  
 Nakata Y, see Ichinose F  
 Namiki A, see Yamakage M  
 Nathalie D, see Eyraud D  
 Neal JM, see Hodgson PS  
 Nellgård B, see Miura Y  
 Nelson JB, Oppenorth TJ, Fleisher LA, Frank SM. Perioperative plasma endothelin-1 and big endothelin-1 concentrations in elderly patients undergoing major surgical procedures, 898  
 Nemirov MA, see Manecke GR Jr  
 Neuwelt EA, see Remsen LG  
 Newman MAJ, see Weightman WM  
 Newton D, see Kochs E  
 Nguyen H, see Talke PO  
 Nguyen JM, see Frank SM  
 Nightingale CE, see Myles PS  
 Nijima K, Seto A, Aoyama K, Takenaka I, Kadoya T. An illuminating stylet as an aid for tracheal intubation via the intubating laryngeal mask airway (letter), 470  
 Niimi Y, see Ichinose F  
 Nikolov MP, see Chaney MA  
 Nishanian EV, Goudsouzian NG. Carbon dioxide embolism in an infant (letter, reply), 227  
 Nishihara M, see Sugiyama K  
 Nishikawa T, see Tanaka M  
 Nishina K, Akamatsu H, Mikawa K, Shiga M, Maekawa N, Obara H, Niwa Y. The effects of clonidine and dexmedetomidine on human neutrophil functions, 452  
 Nishiyama M, Naganuma K, Amaki Y. A

- new approach for brachial plexus block under fluoroscopic guidance, 91
- Nishman RJ, see Rozner MA
- Niwa Y, see Nishina K
- Noe CE, see Ghoname EA
- Nogami WM, see Palmer CM
- Norman GS, see Kelly DJ
- Noronha D, see Lauwers M
- Norris MC, Ferrenbach D, Dalman H, Fogel ST, Borrenpohl S, Hoppe W, Riley A. Does epinephrine improve the diagnostic accuracy of aspiration during labor epidural analgesia?, 1073
- see Huffnagle S
- Nun MB, see Fredman B
- Nyström EUM, Heavner JE, Buffington CW. Blood pressure is maintained despite profound myocardial depression during acute bupivacaine overdose in pigs, 1143
- Obara H, see Nishina K
- Oczenski W, Krenn H, Dahaba AA, Binder M, El-Schahawi-Kienzl I, Jellinek H, Schwarz S, Fitzgerald RD. Hemodynamic and catecholamine stress responses to insertion of the Combitube, laryngeal mask airway or tracheal intubation, 1389
- Odonkor P, see Fahy BG
- Ogawa R, see Shiga T
- Ogura T, see Otsuka F
- Oh Y. Peripheral Nerve Blocks, Upper and Lower Limb (book review), 697
- Ohmura S, Ohta T, Yamamoto K, Kobayashi T. A comparison of the effects of propofol and sevoflurane on the systemic toxicity of intravenous bupivacaine in rats, 155
- Ohnishi Y, see Inoue S
- Ohta T, see Ohmura S
- Ohta Y, see Kanda T
- Oka K, see Furuya R
- Okada K, see Fujioka S
- Ökten F see Kurtipek Ö
- Okubo N, see Bouaziz H
- Okumura F, see Soga K
- Olçay E, see Celebioglu B
- Olinger A, see Vollmar B
- Olmedilla L, see Garutti I
- Olufolabi Y, see Gan TJ
- Omar A, see Abderrahmane B
- Ong BC, see Chow MYH
- Ongerboer de Visser BW, see Ubags LH
- Ono Y, see Fujioka S
- Opgenorth TJ, see Nelson JB
- Oral M, see Kurtipek Ö
- O'Reilly M, see Cornish PB
- Ornaque I, see Suarez S
- Orszulak TA, see Plöchl W
- Oshita S, see Kawahito S
- Österlund Å, Arlander E, Eriksson LJ, Lindahl SGE. The effects on resting ventilation of intravenous infusions of morphine or sameridine, a novel molecule with both local anesthetic and opioid properties, 160
- Otsuka F, Morita K, Takeuchi M, Yamauchi T, Ogura T, Sekines K, Miura M, Hirakawa M, Makino H. The effects of intrinsic vasopressin on urinary aquaporin-2 excretion and urine osmolality during surgery under general anesthesia, 181
- Ovali E, see Doğan IV
- Overdyk FJ, Harvey SC. Successful strategies for improving operating room efficiency at academic institutions (letter, reply), 963
- Owen-Reece H, see Lovell AT
- Paech M, see Scott DA
- Pagel MA, see Remsen LG
- Palitzsch K-D, see Keyl C
- Palmer CM, Maciulla JE, Cork RC, Nogami WM, Gossler K, Alves D. The incidence of fetal heart rate changes after intrathecal fentanyl labor analgesia, 577
- Palmer PP. Implantable pain therapies—what training is required?, 1203
- Pan M-H, see Cheng J-K
- Pancrazio JJ, see McDowell TS
- Pandey CK, see Lakhotia R
- Pandey JC, see Sinha PK
- Pandit JJ. Hypoxic ventilatory response: the effects of CO<sub>2</sub> and of sustained hypoxia (letter), 695
- Pandit UA, see Tait AR
- Pannetier J-C, see Gueugniaud P-Y
- Panning B, see Walz R
- Parakkamodom S, see Visalyaputra S
- Pardo H, see Janiszewski DJ
- Parlow JL, Sagnard P, Begou G, Viale J-P, Quintin L. The effects of clonidine on sensitivity to phenylephrine and nitroprusside in patients with essential hypertension recovering from surgery, 1239
- Patel PM, see Soonthon-Brant V
- Pearlstein RD
- see Grocott HP
- see Miura Y
- Peter K, see Gries A
- Peterson LR, see Vidovich MI
- Pethpaisit N, see Visalyaputra S
- Phillips J. Ondansetron versus droperidol for postoperative nausea and vomiting (letter), 465
- Phillips-Bute B, see Gan TJ
- Picker MJ, see Morgan D
- Piegras DG, see Kuhnert SM
- Pietrabissa R, see Runza M
- Piper S, see Suttner S
- Pitkänen M, see Heavner JE
- Plöchl W, Cook DJ, Orszulak TA, Daly RC. Intracranial pressure and venous cannulation for cardiopulmonary bypass, 329
- Plotkin JS, see Fahy BG
- Pochettino A, see Cheung AT
- Pogatzki E, see Brodner G
- Pohorille A, see Eger EI II
- Polissar NL, see Domino KB
- Politis GD, Tobin JR, Morell RC, James RL, Cantwell MF. Tracheal intubation of healthy pediatric patients without muscle relaxant: a survey of technique utilization and perceptions of safety, 737
- Pollock JE, see Hodgson PS
- Polosa C, see Backman SB
- Ponsford JL, see Myles PS
- Poppers PJ, see Manecke GR Jr
- Posch M, see Knapp S
- Posner KL, see Domino KB
- Power I, see Asai T
- Prichard CB, see Greenberg RS
- Primorac D, see Kapural L
- Prohaska B, see Kapral S
- Proost JH, see Schiere S
- Ptacek LJ, see Viscomi CM
- Pühringer F, see Keller C
- Pulido C, see Ferrando A
- Puura AIE, Rorarius MGF, Manninen P, Hoppert S, Baer GA. The costs of intense neuromuscular block for anesthesia during endolaryngeal procedures due to waiting time, 1335
- Quaglini V, see Runza M
- Quintana B, see Garutti I
- Quintin L, see Parlow JL
- Quirion R, see Gilron I
- Raby KE, Brull SJ, Timimi F, Akhtar S, Rosenbaum S, Naimi C, Whittemore AD. The effect of heart rate control on myocardial ischemia among high-risk patients after vascular surgery, 477
- Rädler C, see Keller C
- Radzievsky AA, Rojavin M, Cowan A, Ziskin MC. Suppression of pain sensation caused by millimeter waves: a double-blinded, cross-over, prospective human volunteer study, 836
- Rajek A, Lenhardt R, Sessler DI,

- Grabenwöger M, Kastner J, Mares P, Jantsch U, Gruber E. Tissue heat content and distribution during and after cardiopulmonary bypass at 17°C, 1220
- Ralley FE. The use of nitric oxide for managing catastrophic pulmonary vasoconstriction arising from protamine administration, 505
- Ramanathan J, Angel JJ, Bush AJ, Lawson P, Sibai B. Changes in maternal middle cerebral artery blood flow velocity associated with general anesthesia in severe preeclampsia, 357
- Ramsay MAE. Fat embolism and neurological dysfunction (letter, reply), 1427
- Ramza JT, see Valley RD
- Rasch DK, see Wells LT
- Rassias AJ, Marrin AS, Arruda J, Whalen PK, Beach M, Yeager MP. Insulin infusion improves neutrophil function in diabetic cardiac surgery patients, 1011
- Rattan KN, see Gupta R
- Rehman M, see Shevchenko Y
- Reich DL, Uysal S, Bodian CA, Gabriele S, Hibbard M, Gordon W, Sliwinski M, Kayne RD. The relationship of cognitive, personality, and academic measures to anesthesiology resident clinical performance, 1092
- Reimann W, Schlütz H, Selve N. The antinociceptive effects of morphine, desipramine, and serotonin and their combinations after intrathecal injection in the rat, 141
- Remsen LG, Pagel MA, McCormick CI, Fiamengo SA, Sexton G, Neuwelt EA. The influence of anesthetic choice,  $P_{aCO_2}$ , and other factors on osmotic blood-brain barrier disruption in rats with brain tumor xenografts, 559
- Rennotte MTh, Baele Ph, Aubert G, Rodenstein DO. Epidural opioids and respiratory arrests (letter), 862
- Reuben SS, Connelly NR. Postoperative analgesia for outpatient arthroscopic knee surgery with intraarticular clonidine, 729
- Connelly NR, Maciolek H. Postoperative analgesia with controlled-release oxycodone for outpatient anterior cruciate ligament surgery, 1286
- Steinberg RB, Lurie SD, Gibson CS. A dose-response study of intravenous regional anesthesia with meperidine, 831
- Rich GF, see Locke MM
- Richardson CA, see Talke PO
- Richardson MG, Wissler RN. The effects of needle bevel orientation during epidural catheter insertion in laboring parturients, 352
- Ricksten S-E. The effects of propofol on cerebral blood flow velocity and cerebral oxygen extraction during cardiopulmonary bypass (letter, reply), 232
- see Valsson F
- Riley A, see Norris MC
- Riou B, see Vivien B
- Rodenstein DO, see Rennotte MTh
- Roewer N, see Greim C-A
- Roizen MF. Clinical Anesthesia Procedures of the Massachusetts General Hospital. 5th ed (book review), 236, 472
- Rojavin MA, see Radzievsky AA
- Rolf N, see Loick HM
- Rommelspacher H, see Spies CD
- Rorarius MGF, see Puura AIE
- Rose DK, see Joo HS
- Rose JB, Cuy R, Cohen DE, Schreiner MS. Preoperative oral dextromethorphan does not reduce pain or analgesic consumption in children after adenotonsillectomy, 749
- Rose RJ, see Fanciullo GJ
- Rosen DA, see Valenzuela RC
- Rosen M, see Lauwers M
- Rosenbaum S, see Raby KE
- Rosenberg P, see Heavner JE
- Ross E, see Fanciullo GJ
- Roth SH, see Wakasugi M
- Rothschiller JL, Uejima T, Dsida RM, Côté CJ. Evaluation of a new operating room ventilator with volume-controlled ventilation: the Ohmeda 7900, 39
- Røttnes JS, see Klaastad Ø
- Rouine-Rapp K, see Lutz JT
- Rowlingson J. Epidural steroids in treating failed back surgery syndrome (editorial), 240
- Roytblat L, see Talmor D
- Rozner MA, Nishman RJ. Pacemaker-driven tachycardia revisited (letter), 965
- Rubinstein A, see Myles PS
- Rudow MP, see Freedman JM
- Runza M, Pietrabissa R, Mantero S, Albani A, Quaglini V, Contro R. Lumbar dura mater biomechanics: experimental characterization and scanning electron microscopy observations, 1317
- Sá Rêgo MM, Inagaki Y, White PF. Remifentanyl administration during monitored anesthesia care: are intermittent boluses an effective alternative to a continuous infusion?, 518
- Inagaki Y, White PF. The cost-effectiveness of methohexital versus propofol for sedation during monitored anesthesia care, 723
- Saatte S, see Khorasani A
- Sagnard P, see Parlow JL
- Saito Y
- see Sakura S
- see Valsson F
- Saitoh Y, see Fujii Y
- Sakabe T, see Wakamatsu H
- Sakaki T, see Kawaguchi M
- Sakamoto A, see Shiga T
- Sakamoto T, see Kawaguchi M
- Sakura S, Sumi M, Kushizaki H, Saito Y, Kosaka Y. Concentration of lidocaine affects intensity of sensory block during lumbar epidural anesthesia, 123
- Samii K, see Bouaziz H
- Samso E, see Farber NE
- Sarantopoulos C, see Fassoulaki A
- Sarraf-Yazdi S, see Grocott HP
- Sasaki T, see Watanabe I
- Sato S, see Ikeda T
- Satoh K, see Sugiyama K
- Satoh T, see Sumida M
- Savage RM, Cosgrove DM. Systematic transesophageal echocardiographic examination in mitral valve repair: the evolution of a discipline into the twenty-first century, 1197
- Savarese JJ, see Lien CA
- Schäfer M, see Brack A
- Schaper A. Rediscovering the obvious (letter), 1427
- Scheld HH, see Loick HM
- Schellenberg AG. Anesthesiology Pocket Guide (book review), 236
- Scheller M, see Schneck H
- Schepens MA, see van Dongen EP
- Schiere S, Proost JH, Schuringa M, Wierda JMKH. Pharmacokinetics and pharmacokinetic-dynamic relationship between rapacuronium (Org 9487) and its 3-desacetyl metabolite (Org 9488), 640
- Schiller NB, see Lambert A-S
- Schlütz H, see Reimann W
- Schmeising C, see Habibi A
- Schmeling WT, Ganjoo P, Staunton M, Drexler C, Farber NE. Pretreatment

- with dexmedetomidine: altered indices of anesthetic depth for halothane in the neuraxis of cats, 625
- see Farber NE
- Schmelz M, see Koppert W
- Schmidt C
- see Loick HM
- see Suttner S
- Schneck H, Scheller M, Wagner R, von Hundelshausen B, Kochs E. Anesthesia for cesarean section and acid aspiration prophylaxis: a German survey, 63
- Schoenwald PK, see Kapural L
- Schreiner MS, see Rose JB
- Schroeder DR
- see Grady RE
- see Horlocker TT
- Schuepfer G, see Konrad C
- Schug SA, see Scott DA
- Schuringa M, see Schiere S
- Schwabe D, see Farber NE
- Schwartz RE, see Shevchenko Y
- Schwarz S, see Oczenski W
- Schweizer A, see Licker M
- Schymroszyk B, see Holst D
- Scott D, see D'Ercole FJ
- Scott DA, Blake D, Buckland M, Etches R, Halliwell R, Marsland C, Merridew G, Murphy D, Paech M, Schug SA, Turner G, Walker S, Huizar K, Gustafsson U. A comparison of epidural ropivacaine infusion alone and in combination with 1, 2, and 4  $\mu\text{g/mL}$  fentanyl for seventy-two hours of postoperative analgesia after major abdominal surgery, 857
- Segawa H, see Miyazaki Y
- Segerdahl M, see Sjölund K-F
- Segura-Vasi AM, Suelto MD, Boudreaux AM. External jugular vein cannulation for central venous access (letter), 692
- Seiler JG III, see Hord AH
- Sekines K, see Otsuka F
- Selve N, see Reimann W
- Semsroth M, see Kozek-Langenecker SA
- Sessler DI. Recommended exposure limits for desflurane and isoflurane (letter, reply), 1424
- Temperature monitoring and management during neuraxial anesthesia (editorial), 243
- see Fleischmann E
- see Hoerauf KH
- see Ikeda T
- see Rajek A
- Seto A, Aoyama K, Takenaka I, Kadoya T. Ventilation difficulties through the intubating laryngeal mask, 1181
- see Niijima K
- Sewell JC, see Eger EI II
- Sexton G, see Remsen LG
- Shaikh S, see Lauwers M
- Shapira Y, see Talmor D
- Shapiro M, see Fredman B
- Sheminant MR, see Weightman WM
- Sheng H, see Grocott HP
- Shevchenko Y, Jocsos JC, McRae VA, Stayer SA, Schwartz RE, Rehman M, Choudhry DK. The use of lidocaine for preventing the withdrawal associated with the injection of rocuronium in children and adolescents, 746
- Shi B, see Heavner JE
- Shi Y-Q, see Sun X-g
- Shichino T, see Miyazaki Y
- Shiga M, see Nishina K
- Shiga T, Sakamoto A, Koizumi K, Ogawa R. Endoscopic thoracic sympathectomy for primary erythromelalgia in the upper extremities, 865
- Shim JC, see Kim KS
- Shingu K
- see Asai T
- see Shinomura T
- Shinomura T, Nakao S, Adachi T, Shingu K. Clonidine inhibits and phorbol acetate activates glutamate release from rat spinal synaptoneurosome, 1401
- Shogase A, Mizutani K, Toyoda Y. Protecting Ohmeda 7900 Ventilator Flow Sensor, 234
- Shore-Lesserson L, Manspeizer HE, DePerio M, Francis S, Vela-Cantos F, Ergin MA. Thromboelastography-guided transfusion algorithm reduces transfusions in complex cardiac surgery, 312
- Sia ATH, Chong JL, Chiu JW. Combination of intrathecal sufentanil 10  $\mu\text{g}$  plus bupivacaine 2.5 mg for labor analgesia: is half the dose enough?, 362
- Sia S, Lepri A. Clonidine administered as an axillary block does not affect postoperative pain when given as the sole analgesic, 1109
- Siao D, see Grenier B
- Sibai B, see Ramanathan J
- Sinatra RS, see Wang A
- Sinclair R, see Mattsson U
- Singh PK, see Sinha PK
- Singh S, see Sinha PK
- Singhal SK, see Gupta R
- Sinha PK, Pandey JC, Dubey PK, Singh PK, Singh S. Cloth band: another unusual cause of difficult endotracheal intubation (letter), 230
- Sittl R, see Koppert W
- Sjögren D, Lindahl SGE, Sollevi A. Hypoxic ventilatory response: the effects of  $\text{CO}_2$  and of sustained hypoxia (letter, reply), 696
- Sjölund K-F, Segerdahl M, Sollevi A. Adenosine reduces secondary hyperalgesia in two human models of cutaneous inflammatory pain, 605
- Sladen RN, see Kapral S
- Slappendel R, Weber EWG, Bugter MLT, Dirksen R. The intensity of preoperative pain is directly correlated with the amount of morphine needed for postoperative analgesia, 146
- Weber EWG, Dirksen R, Gielen MJM, van Limbeek J. Optimization of the dose of intrathecal morphine in total hip surgery: a dose-finding study, 822
- Sleigh JW, Andrzejowski J, Steyn-Ross A, Steyn-Ross M. The Bispectral Index: a measure of depth of sleep?, 659
- Sliwinski M, see Reich DL
- Slogoff S, see Jellish WS
- Smith BL. Use of the laryngeal mask airway in children with upper respiratory tract infections: beware secretions (letter), 693
- Smith CV, see Bucklin BA
- Smith M, see Lovell AT
- Smith MA, see Morgan D
- Smith MP. Drugs, Diseases, and Anesthesia (book review), 472
- Smith MT, Wright AWE, Williams BE, Stuart G, Cramond T. Cerebrospinal fluid and plasma concentrations of morphine, morphine-3-glucuronide, and morphine-6-glucuronide in patients before and after initiation of intracerebroventricular morphine for cancer pain management, 109
- Sneyd R, see Lauwers M
- Soga K, Fujita H, Andoh T, Okumura F. Retinal artery air embolism in dogs: fluorescein angiographic evaluation of effects of hypotension and hemodilution, 1004
- Sollevi A
- see Sjögren D
- see Sjölund K-F
- Somprakit P, see Visalyaputra S

- Somri M, see Gaitini LA
- Song D, Whitten CW, White PF. Use of remifentanyl during anesthetic induction: a comparison with fentanyl in the ambulatory setting, 734  
see White PF
- Sonner JM, see Eger EI II
- Soonthon-Brant V, Patel PM, Drummond JC, Cole DJ, Kelly PJ, Watson M. Fentanyl does not increase brain injury after focal cerebral ischemia in rats, 49
- Sossai R, see Jöhr M
- Spendlove JL, see Girling KJ
- Spiekermann BF, see Locke MM
- Spies CD, Rommelspacher H. Alcohol withdrawal in the surgical patient: prevention and treatment, 946  
see Hansen D
- Spoto M-R, see Laviola S
- Sprung J, see Kapural L
- Stafford M, see Talke PO
- Stamos T, see Hogue CW Jr
- Stanzel M, see Knapp S
- Staunton M  
see Farber NE  
see Schmeling WT
- Stayer SA, see Shevchenko Y
- Stecker MM, see Cheung AT
- Stefanini S, see Varrassi G
- Stein RD, see Backman SB
- Steinberg RB, see Reuben SS
- Stevenson GW, Tobin M, Horn B, Chen EH, Hall SC, Coté CJ. An adult system versus a Bain system: comparative ability to deliver minute ventilation to an infant lung model with pressure-limited ventilation, 527
- Steyn-Ross A, see Sleigh JW
- Steyn-Ross M, see Sleigh JW
- Stienstra R, Dilrosun-Alhadi BZR, Dahan A, van Kleef JW, Veering BT, Burm AGL. The epidural "top-up" in combined spinal-epidural anesthesia: the effect of volume versus dose, 810
- Stockmanns G, see Kochs E
- Stoelting RK, see Gwirtz KH
- Stoiser B, see Knapp S
- Stuart G, see Smith MT
- Su F, see Sun X-g
- Suarez S, Ornaque I, Fábregas N, Valero R, Carrero E. Venous air embolism during Parkinson surgery in patients with spontaneous ventilation, 793
- Suelto MD, see Segura-Vasi A
- Sugiyama K, Yokoyama K, Satoh K, Nishihara M, Yoshitomi T. Does the Murphy eye reduce the reliability of chest auscultation in detecting endobronchial intubation?, 1380
- Sulkes J, see Hochhauser E
- Sullivan LM, Hönenmann CW, Arledge JA, Durieux ME. Synergistic inhibition of lysophosphatidic acid signaling by charged and uncharged local anesthetics, 1117
- Sumi M, see Sakura S
- Sumida M, Inaba H, Isawa E, Fujimoto S, Satoh T, Kawamata M. Prevention by methylprednisolone of increased circulating tumor necrosis factor- $\alpha$  levels and lung injury associated with systemic inflammatory response syndrome due to intraperitoneal hyperthermia, 771
- Summers AC, Gupta AK, Matta BF. Dynamic cerebral autoregulation during sevoflurane anesthesia: a comparison with isoflurane, 341
- Sun X-g, Su F, Shi Y-Q, Lee C. the "second gas effect" is not a valid concept, 188
- Suter PM. Principles and Practice of Intensive Care Monitoring (book review), 967
- Suttner S, Boldt J, Schmidt C, Piper S, Kumle B. Cost analysis of target-controlled infusion-based anesthesia compared with standard anesthesia regimens, 77
- Suwanapeum P, see Visalyaputra S
- Swart M, see Gupta AK
- Syben R, see Hansen D
- Szekely A, Heindl B, Zahler S, Conzen PF, Becker BF. S(+)-ketamine, but not R(-)-ketamine, reduces postischemic adherence of neutrophils in the coronary system of isolated guinea pig hearts, 1017
- Szmuk P, see Ezri T
- Tait AR, Pandit UA, Voepel-Lewis T, Munro HM, Malviya S. Use of the laryngeal mask airway in children with upper respiratory tract infections: beware secretions (letter, reply), 693
- Takenaka I  
see Nijima K  
see Seto A
- Takeuchi M, see Otsuka F
- Talke PO, Caldwell JE, Richardson CA, Kirkegaard-Nielsen H, Stafford M. The effects of dexmedetomidine on neuromuscular blockade in human volunteers, 633
- Tanaka H, see Fujii Y
- Tanaka K, see Kawahito S
- Tanaka M, Nishikawa T. Effective dose of isoproterenol-containing test dose in anesthetized children (letter), 1189
- Nishikawa T. Evaluating T-wave amplitude as a guide for detecting intravascular injection of a test dose in anesthetized children, 754
- Tanil D, see Myles PS
- Tanus-Santos JE, Moreno H Jr. Inhaled nitric oxide and amniotic fluid embolism (letter), 691
- Moreno H Jr, Zappellini A, de Nucci G. Small-dose inhaled nitric oxide attenuates hemodynamic changes after pulmonary air embolism in dogs, 1025  
see Moreno H Jr
- Tarnow P, see Mattsson U
- Tarshis J. Epidural analgesia and cesarean delivery: what is the relationship? (letter), 963
- Taslimi R, see Hoerauf KH
- Taves D, see Webster AC
- ter Beek HT, see van Dongen EP
- Terui K, see Ichinose F
- Tewari P, see Lakhota R
- Thalhammer F, see Knapp S
- Thiel H, see Greim C-A
- Thng CH, see Chow MYH
- Thomas PS. Churchill's Pocketbook of Pain Management (book review), 968
- Hakim TS, Trang LW, Hosain SI, Camporesi EM. The synergistic effect of sympathectomy and hyperbaric oxygen exposure on transcutaneous  $PO_2$  in healthy volunteers, 67
- Thornton C, see Kochs E
- Ti LK, Chow MYH, Lee TL. Comparison of sevoflurane with propofol for laryngeal mask airway insertion in adults, 908
- Tighe KE, Webb AM, Hobbs GJ. Persistently high plasma morphine-6-glucuronide levels despite decreased hourly patient-controlled



- analgesia morphine use after single-dose diclofenac: potential for opioid-related toxicity, 1137
- Tillinger M, see Fredman B
- Timimi F, see Raby KE
- Tinker JH, see Bucklin BA
- Tobias JD, Wilson WR Jr, Meyer DJ.  
Transcutaneous monitoring of carbon dioxide tension after cardiothoracic surgery in infants and children, 531
- Tobin JR, see Politis GD
- Tobin M, see Stevenson GW
- Toh KW, Deacock SJ, Fawcett WJ. Severe anaphylactic reaction to cisatracurium, 462
- Torralva PR, Macario A, Brock-Utne G.  
Another use of a bronchoscopic swivel adapter (letter), 1187
- Torri G  
see Casati A  
see Fanelli G
- Toyoda Y, see Shogase A
- Toyooka H, see Fujii Y
- Tramèr MR, Walder B. Efficacy and adverse effects of prophylactic antiemetics during patient-controlled analgesia therapy: a quantitative systematic review, 1354
- Trang LW, see Thomas PS
- Trudell JR  
see Eger EI II  
see Ueno S
- Truong P, see Macario A
- Tschernko EM, Kritzing M, Gruber EM, Jantsch-Watzinger U, Jandrasits O, Mares P, Wissner W, Klepetko W, Haider W. Lung volume reduction surgery: preoperative functional predictors for postoperative outcome, 28
- Tschopp M, see Konrad C
- Tschuor S, see Laviola S
- Tsen LC, Datta S. Panic attacks and lactated Ringer's solution: is there a relationship?, 795  
see Martin R
- Tsitsika MV, see Mentzelopoulos SD
- Tsujiguchi N, see Yamakage M
- Tsuneyoshi I, see Hamu Y
- Tuman KJ, McCarthy RJ. Individualizing  $\beta$ -adrenergic blocker therapy: patient-specific target-based heart rate control, 475
- Tuovinen K, see Kokki H
- Turner G, see Scott DA
- Tüzün F see Kurtipek Ö
- Twetman ER, see Islander G
- Tzeng G, see Martin R
- Ubags LH, Kalkman CJ, Been HD, Koelman JH, Ongerboer de Visser BW. A comparison of myogenic motor evoked responses to electrical and magnetic transcranial stimulation during nitrous oxide/opioid anesthesia, 568
- Uchida H, see Ichinose F
- Uejima T, see Rothschilder JL
- Ueno S, Trudell JR, Eger EI II, Harris RA.  
Actions of fluorinated alkanols on GABA<sub>A</sub> receptors: relevance to theories of narcosis, 877
- Uetsuki N, see Adachi T
- Uezono S, see Ichinose F
- Utsumi J, see Miyazaki Y
- Uysal S, see Reich DL
- Vaida SJ, see Gaitini LA
- Vakharia AS, see Ghoname EA
- Vakkuri AP, Lindgren L, Korttila KT, Yli-Hankala AM. Transient hyperdynamic response associated with controlled hypocapnic hyperventilation during sevoflurane-nitrous oxide mask induction in adults, 1384
- Valenzuela RC, Rosen DA. Topical lidocaine-prilocaine cream (EMLA) for thoracostomy tube removal, 1107
- Valero R, see Suarez S
- Valley RD, Ramza JT, Calhoun P, Freid EB, Bailey AG, Kopp VJ, Georges LS. Tracheal extubation of deeply anesthetized pediatric patients: a comparison of isoflurane and sevoflurane, 742
- Valsson F, Lundin S, Kirnö K, Hedner T, Houltz E, Saito Y, Ricksten S-E. Atrial natriuretic peptide attenuates pacing-induced myocardial ischemia during general anesthesia in patients with coronary artery disease, 279
- Van Aken H  
see Booke M  
see Brodner G  
see Loick HM
- van Dongen EP, ter Beek HT, Schepens MA, Morshuis WJ, Langemeijer HJ, de Boer A, Boezeman EH. Within-patient variability of myogenic motor-evoked potentials to multipulse transcranial electrical stimulation during two levels of partial neuromuscular blockade in aortic surgery, 22
- van Kleef JW, see Stienstra R
- van Limbeek J, see Slappendel R
- van Meerbeek JP, see Medici G
- van Wezel HB, see Kal JE
- Vandendriessche M, see Grenier B
- Vargas O, see Hansen D
- Varrassi G, Marinangeli F, Agrò F, Aloe L, De Cillis P, De Nicola A, Giunta F, Ischia S, Ballabio M, Steafanini S. A double-blinded evaluation of propacetamol versus ketorolac in combination with patient-controlled analgesia morphine: analgesic efficacy and tolerability after gynecologic surgery, 611
- Veering BT, see Stienstra R
- Veglia F, see Casati A
- Vekeman D, see Goldberg ME
- Vela-Cantos F, see Shore-Lesserson L
- Verchère E, see Grenier B
- Vergroesen I, see Kal JE
- Verhoeven GT, see Medici G
- Viale J-P, see Parlow JL
- Vidne B, see Hochhauser E
- Vidovich MI, Peterson LR, Wong HY. The effect of lidocaine on bacterial growth in propofol, 936
- Vinik HR, Bradley EL Jr, Kissin I. Isobolographic analysis of propofol-thiopental hypnotic interaction in surgical patients, 667
- Visalyaputra S, Lertakyananee J, Pethpaisit N, Somprakit P, Parakkamodom S, Suwanapeum P. Intraperitoneal lidocaine decreases intraoperative pain during postpartum tubal ligation, 1077
- Visco E, see Dunbar PJ
- Viscomi CM, Ptacek LJ, Dudley D. Anesthetic management of familial hypokalemic periodic paralysis during parturition, 1081
- Vivien B, Langeron O, Coriat P, Riou B. Minimum alveolar anesthetic concentration of volatile anesthetics in normal and cardiomyopathic hamsters, 489
- Voepel-Lewis T  
see Huntington JH  
see Tait AR
- Vollmar B, Olinger A, Hildebrandt U, Menger MD. Cardiopulmonary dysfunction during minimally invasive thoraco-lumboendoscopic spine surgery, 1244
- von Hundelshausen B, see Schneck H
- Wachowski I, Jolly DT, Hrazdil J, Galbraith JC, Greacen M, Clanachan AS. The growth of microorganisms in propofol and mixtures of propofol and lidocaine, 209
- Wagner R, see Schneck H
- Wakamatsu H, Matsumoto M,

- Nakakimura K, Sakabe T. The effects of moderate hypothermia and intrathecal tetracaine on glutamate concentrations of intrathecal dialysate and neurologic and histopathologic outcome in transient spinal cord ischemia in rabbits, 56
- Wakasugi M, Hirota K, Roth SH, Ito Y. The effects of general anesthetics on excitatory and inhibitory synaptic transmission in area CA1 of the rat hippocampus *in vitro*, 676
- Wakeling H, see Gan TJ
- Walder B, see Tramèr MR
- Walker S, see Scott DA
- Walker SG, see Gwirtz KH
- Wallner T  
see Fleischmann E  
see Hoerauf KH
- Walsh F, see Kelly DJ
- Waltl B, see Kapral S
- Walz R, Davis S, Panning B. Is the Combitube a useful emergency airway device for anesthesiologists? (letter), 233
- Wang A, Sinatra RS. Epidural anesthesia for cesarean section in a patient with von Hippel-Lindau disease and multiple sclerosis, 1083
- Wang J-J, Ho S-T, Lee S-C, Liu Y-C. A comparison among nalbuphine, meperidine, and placebo for treating postanesthetic shivering, 686
- Wang L, see Hall SR
- Wang SM, see Kain ZN
- Warf P, see Jellich WS
- Warner DS  
see Grocott HP  
see Miura Y
- Watanabe I, Andoh T, Furuya R, Sasaki T, Kamiya Y, Itoh H. Depressant and convulsant barbiturates both inhibit neuronal nicotinic acetylcholine receptors, 1406  
see Furuya R
- Watcha MF, White PF. Cost effectiveness ratio: an often misunderstood term (letter, reply), 1191  
see White PF
- Watson J, see Webster AC
- Watson M, see Soonthon-Brant V
- Watson RS, see Zimmerman AA
- Webb AM, see Tighe KE
- Weber EWG, see Slappendel R
- Webster AC, Morley-Forster PK, Janzen V, Watson J, Dain SL, Taves D, Dantzer D. Anesthesia for intranasal surgery: a comparison between tracheal intubation and the flexible reinforced laryngeal mask airway, 421
- Wedel DJ, Horlocker TT. The *Titanic* is not the best analogy for enoxaparin (letter, reply), 229
- Wei T-T, see Cheng J-K
- Weightman WM, Gibbs NM, Sheminant MR, Whitford EG, Mahon BD, Newman MAJ. Drug therapy before coronary artery surgery: nitrates are independent predictors of mortality and  $\beta$ -adrenergic blockers predict survival, 286
- Weinger M, see Macario A
- Weinstabl C, see Kapral S
- Weiss SJ, see Cheung AT
- Weissman C. Pulmonary function after cardiac and thoracic surgery, 1272
- Wells LT, Rasch DK. Emergence 'delirium' after sevoflurane anesthesia: a paranoid delusion?, 1308
- Welte M, see Hansen D
- Wendt M, see Holst D
- Wennberg R, see Manninen PH
- Whalen PK  
see Fanciullo GJ  
see Rassias AJ
- White PF, Song D. New criteria for fast-tracking after outpatient anesthesia: a comparison with the modified Aldrete's scoring system, 1069
- Watcha MF. Has the use of meta-analysis enhanced our understanding of therapies for postoperative nausea and vomiting?, 1200  
see Ghoname EA  
see Sá Régio MM  
see Song D  
see Watcha MF
- Whitford EG, see Weightman WM
- Whittemore AD, see Raby KE
- Whitten CW, Greilich PE, Ivy R, Burkhardt D, Allison PM. D-dimer formation during cardiac and noncardiac thoracic surgery, 1226  
see Song D
- Wierda JMKH, see Schiere S
- Williams BE, see Smith MT
- Williams EL, Hildebrand KL, McCormick SA, Bedel MJ. The effect of intravenous lactated Ringer's solution versus 0.9% sodium chloride solution on serum osmolality in human volunteers, 999
- Williams JK, see Zimmerman AA
- Wilson WR Jr, see Tobias JD
- Winters KJ, see Hogue CW Jr
- Wisser W, see Tschernko EM
- Wissler RN, see Richardson MG
- Wong C-S, see Wu C-T
- Wong HY, see Vidovich MI
- Wright AWE, see Smith MT
- Wu C-T, Yu J-C, Yeh C-C, Liu S-T, Li C-Y, Ho S-T, Wong C-S. Preincisional dextromethorphan treatment decreases postoperative pain and opioid requirement after laparoscopic cholecystectomy, 1331
- Wu K-H, see Cheng J-K
- Willenweber J, see Booke M
- Wysocki TT  
see Cassidy JF Jr  
see Miller KM
- Xue FS, Li BW, Zhang GS, Liao X, Zhang YM, Liu JH, An G, Luo LK. The influence of surgical sites on early postoperative hypoxemia in adults undergoing elective surgery, 213
- Yamakage M, Matsuzaki T, Tsujiguchi N, Mori T, Namiki A. The direct effects of heparin and protamine on canine tracheal smooth muscle tone, 1232
- Yamamoto K, see Ohmura S
- Yamauchi T, see Otsuka F
- Yasui O, see Adachi T
- Yayci A, see Doğan IV
- Yeager MP, see Rassias AJ
- Yeh C-C, see Wu C-T
- Yli-Hankala AM, see Vakkuri AP
- Yokoyama K, see Sugiyama K
- Yoshimura N, see Hamu Y
- Yoshitomi T, see Sugiyama K
- Young JV, see Gwirtz KH
- Yu J-C, see Wu C-T
- Zacny JP, see Janiszewski DJ
- Zafeiridis A, see Goldberg ME
- Zahler S, see Szekely A
- Zahn J, see Beilin Y
- Zappellini A, see Tanus-Santos JE
- Zaugg M, Lucchinetti E, Zollinger A. Is continuous intraarterial blood gas monitoring reliable during one-lung ventilation? (letter, reply), 1192
- Zeck S, see Koppert W
- Zhang GS, see Xue FS
- Zhang YM, see Xue FS
- Zhou W, see Landesburg G
- Zimmerman AA, Watson RS, Williams JK. Protein S deficiency presenting as an acute postoperative arterial thrombosis in a four-year-old child, 535
- Zimpfer M, see Fleischmann E
- Ziskin MC, see Radzievsky AA
- Zohar E, see Fredman B
- Zollinger A, see Zaugg M
- Zolotarsky V, see Hochhauser E
- Zotou M, see Fassoulaki A

## Subject Index

- Abdominal surgery**, see Surgery
- Acid aspiration syndrome**, see Complications
- Acid-base balance**  
saline vs lactated Ringer's, human volunteers, 88:999  
temperature correction, 88:538
- Acupressure**, see Anesthetic techniques
- Acupuncture**, see Anesthetic techniques
- Acute myocardial infarction**, see Complications
- Adenosine**, see Pharmacology
- Adenotonsillectomy**, see Surgery
- Adrenalectomy**, see Surgery
- $\alpha$ -Adrenergic receptors**, see Receptors
- $\beta$ -Adrenergic receptors**, see Receptors
- Adrenomedullin**, see Brain; Peptides, vasodilating
- Age factors**, effects of epidural lidocaine, with and without epinephrine, on spinal cord blood flow, anesthetized rabbits, 88:1302
- Airway**  
complications  
disposable vs reusable LMA, 88:1418  
postoperative macroglossia, rapid resolution after extubation of trachea, 88:220  
device, emergency, for anesthesiologists, use of Combitube (letter), 88:233  
difficult  
management, with balloon inflation (letter), 88:1425  
unanticipated, airway adjunct (letter), 88:467  
management  
difficult airway management with balloon inflation (letter), 88:1425  
repeated use of cuffed oropharyngeal airway for radiation therapy, infant (letter), 88:1421  
pressure, endobronchial intubation, as cause of immediate increase in peak inflation pressure, pediatric patients, 88:268  
sealing and cuff pressure, LMA devices, 88:913  
technique, disposable vs reusable LMA, 88:1418  
tracheal extubation of deeply anesthetized pediatric patients, isoflurane vs sevoflurane, 88:742  
tracheostomy, followed by tracheo-innominate artery fistula, 88:777
- Airway obstruction**, upper, jet ventilation, rediscovering the obvious (letter), 88:1427
- Alcohol withdrawal syndrome**, see Complications
- Alfentanil**, see Anesthetics, intravenous
- Allodynia**, see Pain, neuropathic
- Alphaxalone**, see Anesthetics, intravenous
- Ambulatory anesthesia**, see Anesthesia
- Ambulatory surgery**, see Surgery
- Ambulatory surgery centers (ASC)**, impact of rapid anesthetic recovery protocols on ASC staffing, 88:1053
- Amniotic fluid embolism**, see Complications
- Amsterdam Preoperative Anxiety and Information Scale**, see Measurement techniques
- Analgesics**  
clonidine  
effect on activity of neurons in nucleus raphe magnus, rats, 88:459  
intraarticular, postoperative analgesia for outpatient arthroscopic knee surgery, 88:729  
diclofenac, single-dose, followed by persistently high plasma morphine-6-glucuronide levels despite decreased hourly PCA morphine use, 88:1137  
fentanyl  
addition to lidocaine 1.5%, anesthesia for *in vitro* fertilization, 88:523  
epidural ropivacaine infusion alone vs in combination with 1, 2, and 4  $\mu$ g/mL fentanyl, postoperative analgesia after major abdominal surgery, 88:857  
intrathecal, labor analgesia, incidence of fetal heart rate changes, 88:577  
neurotoxicity of spinal drugs, 88:797  
vs remifentanyl, use during anesthetic induction, ambulatory setting, 88:734  
ketoprofen, intravenous, effect on postoperative epidural sufentanil analgesia, children, 88:1036  
ketorolac  
recovery characteristics of sevoflurane and halothane, bilateral myringotomy and pressure equalization tube insertion, preschool-aged children, 88:34  
vs propacetamol, combined with PCA morphine, analgesic efficacy and tolerability after gynecologic surgery, 88:611
- meperidine**, intravenous regional anesthesia, dose-response study, 88:831
- morphine**  
amount needed for postoperative analgesia, and intensity of preoperative pain, direct correlation, 88:146  
antinociceptive effects, role of intrinsic efficiency and stimulus intensity, 88:402  
CSF and plasma concentrations, before and after initiation of intracerebroventricular morphine for cancer pain management, 88:109  
intrathecal, determination of effective dose for pain relief after cesarean delivery, 88:346  
intrathecal, optimization of dose, total hip surgery, 88:822  
intravenous, vs EMLA cream, for pain relief during thoracostomy tube removal, 88:1107  
morphine-6-glucuronide, persistently high plasma levels despite decreased hourly PCA morphine use after single-dose diclofenac, 88:1137  
neurotoxicity of spinal drugs, 88:797  
peripheral antihyperalgesic effect to heat but not mechanical stimulation, healthy volunteers, after ultraviolet-B irradiation, 88:117  
vs sameridine, intravenous infusions, effects on resting ventilation, 88:160  
nalbuphine, for treating postanesthetic shivering, comparison with meperidine and placebo, 88:686
- opioids**  
epidural, and respiratory arrests (letter), 88:962  
intrathecal, for acute postoperative pain, safety and efficacy, 88:599  
N<sub>2</sub>O/opioid anesthesia, comparison of myogenic motor evoked responses to electrical and magnetic transcranial stimulation, 88:568  
 $\mu$ -opioids, and morphine, antinociceptive effects, interactions between, role of intrinsic efficiency and stimulus intensity, 88:402  
PCA, efficacy and adverse effects of prophylactic antiemetics,

**Analgesics—continued**

- quantitative systematic review, 88:1354
- requirement, after laparoscopic cholecystectomy, effect of preincisional dextromethorphan treatment, 88:1331
- oxycodone, controlled-release, for postoperative analgesia after anterior cruciate ligament surgery, 88:1286
- propacetamol, vs ketorolac, combined with PCA morphine, analgesic efficacy and tolerability after gynecologic surgery, 88:611
- remifentanyl
  - administration, during monitored anesthesia care, intermittent boluses vs continuous infusion, 88:518
  - sevoflurane-remifentanyl, vs sevoflurane-N<sub>2</sub>O, trabecular outflow facility and formation rate of aqueous humor, rabbits, 88:781
  - vs fentanyl, use during anesthetic induction, ambulatory setting, 88:734
- samerdine
  - for spinal anesthesia, safety and efficacy, comparative dose-ranging study with lidocaine for inguinal hernia repair, 88:815
  - vs morphine, intravenous infusions, effects on resting ventilation, 88:160
- sufentanil
  - and bupivacaine, spinal, labor analgesia, effect of spinal clonidine, 88:573
  - epidural, postoperative, effect of intravenous ketoprofen, children, 88:1036
  - intrathecal, 10 µg, plus bupivacaine 2.5 mg, for labor analgesia, effectiveness of half dose, 88:362
  - ropivacaine 0.2% plus sufentanil vs bupivacaine 0.175% plus sufentanil, epidural analgesia after abdominal surgery, 88:128
  - tramadol, added to mepivacaine, effect on duration of axillary brachial plexus blockade, 88:853
  - yohimbine, effect on activity of neurons in nucleus raphe magnus, rats, 88:459
- Anaphylaxis**, severe anaphylactic reaction to cisatracurium, 88:462
- Anatomy**, brachial plexus, fluoroscopically guided brachial plexus block, 88:91
- Anesthesia**
  - ambulatory
    - controlled-release oxycodone for anterior cruciate ligament surgery, 88:1286
    - cost of methohexital vs propofol and monitored anesthesia care, 88:723
    - criteria for fast-tracking, 88:1069
    - discharge and complications after ambulatory anesthesia, 88:508
    - early discharge after laparoscopic cholecystectomy, 88:1280
    - impact of rapid anesthetic recovery protocols on ASC staffing, 88:1053
    - intraarticular clonidine, 88:729
    - in vitro* fertilization and spinal anesthesia, 88:518
    - midazolam and hearing ability and audiological reaction time, 88:1064
    - remifentanyl intermittent boluses vs continuous infusion during MAC, 88:518
    - remifentanyl vs fentanyl and anesthetic induction, 88:734
  - cardiovascular
    - acute myocardial infarction during lung volume reduction surgery, 88:332
    - adrenomedullin and hemorrhagic shock, 88:326
    - aprotinin and cerebral ischemia, 88:1
    - asymmetric retractors and nerve injury, 88:292
    - atrial natriuretic peptide attenuates myocardial ischemia, 88:279
    - autonomic dysfunction and anesthetic induction, 88:985
    - bronchial rupture by double-lumen endobronchial tube, 88:1252
    - cerebral production of adrenomedullin, 88:1030
    - changing temperature management for CPB, 88:1254
    - clonidine effects on sensitivity to phenylephrine and nitroprusside, 88:1239
    - colloids, crystalloids, and coagulation, 88:483
    - D-dimer formation during thoracic surgery, 88:1226
    - drugs and hospital mortality, 88:286
    - heart rate control and ischemia, 88:477
    - heparin and protamine effects on tracheal smooth muscle, 88:1232
    - Hextend plasma expander for major surgery, 88:985
    - insulin and neutrophil function, 88:1011
    - intracranial pressure and venous cannulation, CPB, 88:329
  - intravenous magnesium and platelet function, 88:1213
  - isoflurane, nitroprusside, and protamine interactions, 88:710
  - ketamine stereoisomers and PMN adhesion, 88:1017
  - lung protection during endobronchial laser surgery, 88:298
  - MAC in cardiomyopathic hamsters, 88:489
  - minimally invasive thoraco-lumboendoscopic spine surgery, 88:1244
  - mitral valve regurgitation, TEE examination, 88:1205
  - nitroglycerin and metabolic coronary flow control, 88:271
  - NO: tetracaine and lidocaine cardiac toxicity, 88:717
  - NO and air embolism, 88:1025
  - NO for management of catastrophic pulmonary vasoconstriction, 88:505
  - one-lung ventilation and thoracic epidural, 88:494
  - osmolality: saline vs lactated Ringer's, 88:999
  - pheochromocytoma and laparoscopy, 88:16
  - predictors for lung volume reduction surgery, 88:28
  - pulmonary function after cardiac and thoracic surgery, 88:1272
  - retinal artery air embolism, 88:1004
  - retrograde cerebral perfusion, 88:8
  - selecting size of double-lumen endobronchial tubes, 88:302
  - sympathetic innervation of forelimb, 88:320
  - tachycardia-induced subendocardial necrosis, 88:973
  - TEE and laparoscopic nephrectomy monitoring, 88:500
  - terlipressin and refractory hypotension, 88:980
  - thoracic epidural anesthesia, clonidine, and CABG, 88:701
  - thromboelastography, cardiac surgery, 88:312
  - tissue heat content and distribution during and after CPB, 88:1220
  - transnasal TEE, 88:302
  - variability motor-evoked responses and aortic surgery, 88:22
- critical care and trauma
  - analgesia for thoracostomy tube removal, 88:1107
  - assessment of tracheal tube position, 88:766

- decreased tumor necrosis factor- $\alpha$  and inflammatory syndrome by steroids and intraperitoneal hyperthermia perfusion, 88:771
- effects of vasopressin on human arteries, 88:542
- hypothermia and intrapulmonary gas exchange, 88:538
- sympathectomy and hyperbaric O<sub>2</sub> therapy, 88:67
- tracheo-innominate artery fistula and tracheostomy, 88:777
- economics and health systems research consensus on anesthesia outcomes, 88:1085
- intense neuromuscular block during endolaryngeal procedures, 88:1335
- predictive evaluation of residents, 88:1092
- neurosurgical
- anesthetics, catecholamines, and cerebral ischemia, 88:787
  - brain tissue oxygenation vs SjvO<sub>2</sub> in head injury, 88:549
  - CO<sub>2</sub> gradient and surgical position, 88:43
  - electrical and magnetic transcutaneous stimulation, 88:568
  - fentanyl and focal cerebral ischemia, 88:49
  - localization of epileptogenic focus, 88:1101
  - measurement of cerebral oxygenation, 88:554
  - optimizing blood-brain barrier disruption, 88:559
  - postoperative neurosurgical pain, 88:335
  - sevoflurane, remifentanyl, and aqueous humor dynamics, 88:781
  - sevoflurane and dynamic autoregulation, 88:341
  - tetracaine and spinal cord ischemia, 88:56
  - venous air embolism, Parkinson surgery, patients with spontaneous ventilation, 88:793
- obstetric
- cesarean delivery of twin gestation under 2 MAC isoflurane, one normal and one with large neck mass, 88:584
  - cesarean section, aspiration prophylaxis in Germany, 88:63
  - epidural anesthesia, patient with von Hippel-Lindau disease, 88:1083
  - familial hypokalemic periodic paralysis during parturition, 88:1081
  - granisetron/dexamethasone for PONV, 88:1340
  - intraperitoneal lidocaine during postpartum tubal ligation, 88:1077
  - intrathecal fentanyl and fetal heart rate, 88:577
  - lateral vs cephalad Tuohy bevel orientation, 88:352
  - middle cerebral artery flow velocity, anesthesia, and preeclampsia, 88:357
  - panic attacks and lactated Ringer's solution, relationship, 88:795
  - paravertebral blockade for modified radical mastectomy, pregnant patient, 88:1340
  - pneumocephalus with intense headache and unilateral pupillary dilatation after accidental dural puncture, 88:582
  - reduced dose combined spinal-epidural in labor, 88:362
  - ropivacaine for initiation of labor epidural analgesia, 88:1340
  - spinal clonidine prolongs labor analgesia, 88:573
  - spinal morphine and post-cesarean section analgesia, 88:346
  - testing epidural catheters, 88:1073
- pediatric
- adult circle system use, infant model, 88:527
  - anesthesia and protein S deficiency, 88:535
  - anesthesia induction, behavioral changes, 88:1042
  - carbon dioxide embolism, infant (letter), 88:227
  - cardiac tumor resection, intraoperative management, 88:1048
  - colonic puncture during ilioinguinal nerve block, 88:1051
  - dextromethorphan premedication, 88:749
  - effects of epidural lidocaine on spinal cord blood flow, 88:1302
  - emergence "delirium" after sevoflurane, 88:1308
  - endobronchial intubation, 88:268
  - endobronchial intubation (letter), 88:229
  - evaluation of Ohmeda 7900 ventilator, 88:39
  - granisetron vs perphenazine, PONV after tonsillectomy, 88:1298
  - halothane and heart sounds, 88:263
  - intraoperative autotransfusion, surgery, 88:763
  - intravenous ketoprofen, 88:1036
  - pain on injection of rocuronium, 88:746
  - parental knowledge and anxiety, 88:251
  - patients with malignant osteopetrosis, 88:1292
  - perioperative information and parental anxiety (editorial), 88:237
  - platelet hyporeactivity in young infants, 88:258
  - preanesthetic videotape for parental preparation, 88:246
  - rate of rise of halothane, children with shunt, 88:759
  - recovery from halothane and sevoflurane, 88:34
  - simple PASS (letter), 88:227
  - test dose efficacy, 88:754
  - tracheal extubation during deep anesthesia, 88:742
  - tracheal intubation without muscle relaxants, 88:737
  - transcutaneous CO<sub>2</sub> and cardiac surgery, 88:531
- regional anesthesia and pain management
- activation of rostral ventrolateral medulla during strychnine allodynia, 88:1125
  - adenosine reduces secondary hyperalgesia, 88:605
  - anesthetics and spinal Fos, 88:407
  - assessing level of sensory block, 88:398
  - axillary block: tramadol plus mepivacaine, 88:853
  - balanced postoperative epidural analgesia, 88:128
  - bupivacaine toxicity, propofol, and sevoflurane, 88:155
  - clonidine for brachial plexus block, 88:1109
  - clonidine plus lidocaine for tourniquet pain during intravenous regional anesthesia, 88:1327
  - diclofenac and renal function, 88:149
  - diclofenac during morphine PCA, 88:1137
  - endoscopic thoracic sympathectomy for primary erythromelalgia, upper extremities, 88:865
  - epidural ropivacaine and fentanyl infusion, 88:857
  - estimating brachial plexus depth, 88:1113
  - fluoroscopically guided brachial plexus block, 88:91
  - fluoroscopy and epidural steroids, 88:367



**Anesthesia—continued**

- frequency of electrical stimulation and percutaneous nerve stimulation therapy, 88:841
- hemodynamic effects of epidural phenylephrine, 88:1322
- implantable pain therapies, nationwide survey, 88:1311
- interactions among  $\mu$ -opioids, 88:407
- interscalene block,  $\beta$ -blockers, and vasovagal reactions, 88:378
- interscalene regional anesthesia, prevention of autonomic hyperreflexia, quadriplegic patient undergoing shoulder surgery, 88:98
- intrathecal opioid analgesia for postoperative pain, 88:599
- lidocaine concentration and epidural anesthesia, 88:123
- local anesthetic site of action on LPA signaling, 88:1117
- lumbar dura mater biomechanics, 88:1317
- mechanism of spinal epidural anesthesia, 88:810
- meperidine intravenous regional anesthesia, 88:831
- metal toxicity and combined spinal-epidural anesthesia, 88:393
- metoclopramide, peridural block, and histology, 88:100
- morphine, morphine-3-glucuronide, and morphine-6-glucuronide, in CSF and plasma, 88:109
- morphine and monoamine antinociception, 88:141
- MRI and Raj's infraclavicular block, 88:593
- nerve stimulator for peripheral regional anesthesia, 88:847
- neurologic complications from CSF drains, 88:388
- neurotoxicity of spinal drugs, 88:797
- optimization of intrathecal morphine dose, 88:822
- pain suppression by millimeter waves, 88:836
- patient with postdural puncture headache and acute leukemia, 88:166
- peripheral antihyperalgesia effect of morphine, 88:117
- persistent paresthesia and repeated axillary block, 88:382
- postoperative propacetamol vs ketorolac, 88:611
- preincisional dextromethorphan, laparoscopic cholecystectomy, 88:1331

- preoperative pain enhances postoperative morphine intake, 88:146
- pulmonary changes after interscalene block, 88:587
- rat paw perfusion after mononeuropathy, 88:103
- remifentanyl during regional anesthesia, 88:134
- samerdine and spinal anesthesia, 88:815
- samerdine vs morphine, effects on ventilation, 88:160
- temperature monitoring/regional anesthesia, 88:373
- thoracic epidural anesthesia and visceral perfusion and metabolism, 88:402
- topical local anesthetics in human burns, 88:1131
- warning signs of bupivacaine overdose, 88:1143

**Anesthesia machines, see Equipment****Anesthesiologists**

- expert, panel, and clinical anesthesia outcomes, 88:1085
- exposure to sevoflurane and  $N_2O$ , 88:925
- use of Combitube as emergency airway device (letter), 88:233

**Anesthetic delivery systems, see Equipment****Anesthetics, gases**

- nitrous oxide ( $N_2O$ )
  - combined with sevoflurane, subanesthetic concentrations, effect on analgesia, mood, and psychomotor performance, healthy volunteers, 88:1149
- exposure, during four different methods of anesthetic induction, 88:925
- fentanyl/ $N_2O$ , effects on concentrations of brain and plasma catecholamines during near-complete cerebral ischemia, rats, 88:787
- $N_2O$ /opioid anesthesia, comparison of myogenic motor evoked responses to electrical and magnetic transcranial stimulation, 88:568
- 1 MAC isoflurane/ $N_2O$  anesthesia, movement to noxious stimulation during, prediction, middle latency auditory evoked responses and EEG-derived variables, 88:114288:1412
- "second gas effect" is not valid concept, 88:188

- sevoflurane- $N_2O$ , mask induction, transient hyperdynamic response associated with controlled hypocapnic hyperventilation during, adults, 88:1384
- sevoflurane- $N_2O$ , vs sevoflurane-remifentanyl, trabecular outflow facility and formation rate of aqueous humor, rabbits, 88:781
- subanesthetic concentrations, effect on analgesia, mood, and psychomotor performance, healthy volunteers, 88:1149
- variability motor-evoked responses and aortic surgery, 88:22
- vs xenon, inhibitory effects on spinal dorsal horn neurons, spinal cord transected cats, 88:884
- xenon, vs  $N_2O$ , inhibitory effects on spinal dorsal horn neurons, spinal cord transected cats, 88:884

**Anesthetics, intravenous**

- alfentanil, intraoperative localization of epileptogenic focus, 88:1101
- alphaxalone, pre- vs postinjury effects on formalin-induced Fos immunoreactivity, spinal cord, rats, 88:414
- dexmedetomidine
  - effects on neuromuscular blockade, human volunteers, 88:633
- modulation of cardiovascular responses to stimulation of CNS pressor sites, 88:617
- pretreatment with, altered indices of anesthetic depth for halothane, neuraxis of cats, 88:625
- etomidate, measurement of cerebral oxygenation, 88:554
- fentanyl
  - and brain injury after focal cerebral ischemia, rats, 88:49
- fentanyl/ $N_2O$ , effects on concentrations of brain and plasma catecholamines during near-complete cerebral ischemia, rats, 88:787
- intraoperative localization of epileptogenic focus, 88:1101
- fluorinated alkanols, MAC, relevance to theories of narcosis, rats, 88:867
- ketamine
  - effects on concentrations of brain and plasma catecholamines during near-complete cerebral ischemia, rats, 88:787
  - effects on excitatory and inhibitory synaptic transmission, area CA1 of rat hippocampus *in vitro*, 88:676

- effects on neuronal nicotinic acetylcholine receptors and  $P_{2X}$  purinoceptors in PC12 cells, 88:174
- isomers, S(+)-ketamine vs R(-)-ketamine, effect on postischemic adherence of neutrophils, coronary system of isolated guinea pig hearts, 88:1017
- neurotoxicity of spinal drugs, 88:797
- meperidine, for treating postanesthetic shivering, comparison with meperidine and placebo, 88:686
- methohexital, vs propofol, cost-effectiveness for sedation during monitored anesthesia care, 88:723
- midazolam
- effects on pure tone audiometry, speech audiometry, and audiological reaction times, human volunteers, 88:1064
  - safety and effectiveness of remifentanyl as adjunct sedative for regional anesthesia, 88:134
- pentobarbital
- effects on excitatory and inhibitory synaptic transmission, area CA1 of rat hippocampus *in vitro*, 88:676
  - pre- vs postinjury effects on formalin-induced Fos immunoreactivity, spinal cord, rats, 88:414
- propofol
- bacterial growth, effect of lidocaine, 88:936
  - effects on cerebral blood flow velocity and cerebral oxygen extraction during CPB (letter), 88:231
  - effects on excitatory and inhibitory synaptic transmission, area CA1 of rat hippocampus *in vitro*, 88:676
  - effects on neuronal nicotinic acetylcholine receptors and  $P_{2X}$  purinoceptors in PC12 cells, 88:174
  - in vitro* effects on platelet aggregation, 88:432
  - growth of microorganisms, 88:209
  - and lidocaine, mixtures, growth of microorganisms, 88:209
  - for lidocaine-induced seizures (letter), 88:1193
  - measurement of cerebral oxygenation, 88:554
  - pre- vs postinjury effects on formalin-induced Fos immunoreactivity, spinal cord, rats, 88:414
  - target plasma concentration required to place LMA vs cuffed oropharyngeal airway, 88:917
  - thiopental hypnotic interaction, isobolographic analysis, surgical patients, 88:667
  - variability motor-evoked responses and aortic surgery, 88:22
  - vs methohexital, cost-effectiveness for sedation during monitored anesthesia care, 88:723
  - vs sevoflurane, core hypothermia, 88:921
  - vs sevoflurane, effects on systemic toxicity of intravenous bupivacaine, rats, 88:155
  - vs sevoflurane, LMA insertion, adults, 88:908
- remifentanyl, as adjunct sedative for regional anesthesia, safety and effectiveness, 88:134
- thiopental
- measurement of cerebral oxygenation, 88:554
  - propofol hypnotic interaction, isobolographic analysis, surgical patients, 88:667
- Anesthetics, local**
- benzocaine, ointment containing, as cause of suspected severe methemoglobinemia (letter), 88:1190
- bupivacaine
- bupivacaine 2.5 mg plus intrathecal sufentanil 10  $\mu$ g, for labor analgesia, effectiveness of half dose, 88:362
  - bupivacaine 0.175% plus sufentanil, vs ropivacaine 0.2% plus sufentanil, epidural analgesia after abdominal surgery, 88:128
  - clonidine and, intraarticular administration, postoperative analgesia for outpatient arthroscopic knee surgery, 88:729
  - intravenous, systemic toxicity, comparison of effects of propofol and sevoflurane, rats, 88:155
  - nerve stimulator for peripheral regional anesthesia, 88:847
  - neurotoxicity of spinal drugs, 88:797
  - overdose, acute, maintenance of blood pressure during profound myocardial depression, pigs, 88:1143
  - sufentanil and, spinal, labor analgesia, effect of spinal clonidine, 88:573
  - charged and uncharged, synergistic inhibition of lysophosphatidic acid signaling, 88:1117
- EMLA cream
- postburn, effect, digital image analysis of erythema development after experimental thermal injury to human skin, 88:1131
- vs intravenous morphine, for pain relief during thoracostomy tube removal, 88:1107
- lidocaine
- assessing level of sensory block, 88:398
  - cardiotoxicity, modification by nitric oxide synthesis inhibition, 88:717
  - concentration, effect on intensity of sensory block during lumbar epidural anesthesia, 88:123
  - effect on bacterial growth in propofol, 88:936
  - epidural, followed by bilateral buttock and leg pain (letter), 88:1188
  - epidural, with and without epinephrine, age-related effects on spinal cord blood flow, anesthetized rabbits, 88:1302
  - induced seizures, treatment with propofol (letter), 88:1193
  - intraperitoneal, and intraoperative pain during postpartum tubal ligation, 88:1077
  - lidocaine 1.5%, addition of fentanyl to, anesthesia for *in vitro* fertilization, 88:523
  - nerve stimulator for peripheral regional anesthesia, 88:847
  - neurotoxicity of spinal drugs, 88:797
  - plus clonidine, for intravenous regional anesthesia, prevention of tourniquet pain, 88:1327
  - propofol and, mixtures, growth of microorganisms, 88:209
  - use for preventing withdrawal associated with injection of rocuronium, children and adolescents, 88:746
  - vs sameridine, safety and efficacy for spinal anesthesia, 88:815
- meperidine, for spinal anesthesia, effects of three graded doses, African men, 88:827
- mepivacaine
- nerve stimulator for peripheral regional anesthesia, 88:847
  - reversal of prolonged unconsciousness by naloxone after intravascular injection of local anesthetic and clonidine, 88:1185
  - tramadol added to, effect on duration of axillary brachial plexus blockade, 88:853
  - 2%, vs 0.5% and 0.75% ropivacaine, interscalene brachial plexus block, pulmonary function changes, double-blinded comparison, 88:587

**Anesthetics, local—continued****ropivacaine**

- epidural, for initiation of labor
  - epidural analgesia, dose-finding study, 88:1340
- epidural infusion, alone vs in combination with 1, 2, and 4  $\mu$ g/mL fentanyl, postoperative analgesia after major abdominal surgery, 88:857
- reversal of prolonged unconsciousness by naloxone after intravascular injection of local anesthetic and clonidine, 88:1185
- ropivacaine 0.2% plus sufentanil, vs bupivacaine 0.175% plus sufentanil, epidural analgesia after abdominal surgery, 88:128
- vs mepivacaine, interscalene brachial plexus block with 0.5% and 0.75% ropivacaine, pulmonary function changes, double-blinded comparison with 2% mepivacaine, 88:587

**samerdine**

- for spinal anesthesia, safety and efficacy, comparative dose-ranging study with lidocaine for inguinal hernia repair, 88:815
- vs morphine, intravenous infusions, effects on resting ventilation, 88:160

**tetracaine**

- cardiotoxicity, modification by nitric oxide synthesis inhibition, 88:717
- transient spinal cord ischemia and, rabbits, 88:56

**Anesthetics, volatile****desflurane**

- confusion, and sevoflurane (letter), 88:1194
- MAC, normal and cardiomyopathic hamsters, 88:489
- recommended exposure limits (letter), 88:1424
- vaporizers, use of electricity (letter), 88:1189

**enflurane**

- effects on depolarization-evoked glutamate release, rat cortical brain slices, 88:1168
- "second gas effect" is not valid concept, 88:188
- sensitivity of T-type calcium currents in various cell types, 88:168
- fluorinated alkanols, actions on GABA<sub>A</sub> receptors, relevance to theories of narcosis, 88:877

**halothane**

- anesthetic depth, and pretreatment with dexmedetomidine, cats, 88:625
- arterial and end-tidal, rate of rise, effect of right-to-left intracardiac shunt, children, 88:759
- combined with dexmedetomidine, and CNS pressor sites, 88:617
- effect on amplitude and frequency characteristics of heart sounds, children, 88:263
- effects on excitatory and inhibitory synaptic transmission, area CA1 of rat hippocampus *in vitro*, 88:676
- lipid emulsion (5% vol/vol), intravenous administration, anesthetic and physiologic effects, 88:671
- MAC, normal and cardiomyopathic hamsters, 88:489
- recovery characteristics, preschool-aged children undergoing bilateral myringotomy and pressure equalization tube insertion, 88:34
- sensitivity of T-type calcium currents in various cell types, 88:168
- hypothesis, production of immobility by acting on two sites approximately five carbon atoms apart, 88:1395

**isoflurane**

- effect on motoneuron excitability by direct spinal action, F-wave study, 88:681
- effects on concentrations of brain and plasma catecholamines during near-complete cerebral ischemia, rats, 88:787
- effects on depolarization-evoked glutamate release, rat cortical brain slices, 88:1168
- effects on excitatory and inhibitory synaptic transmission, area CA1 of rat hippocampus *in vitro*, 88:676
- fentanyl and focal cerebral ischemia, rats, 88:49
- in vitro* effects on platelet aggregation, 88:432
- MAC, normal and cardiomyopathic hamsters, 88:489
- MAC, 1, N<sub>2</sub>O anesthesia, movement to noxious stimulation during, prediction, middle latency auditory evoked responses and EEG-derived variables, 88:114288:1412
- MAC, 2, cesarean delivery of twin gestation, one normal and one with large neck mass, 88:584
- recommended exposure limits (letter), 88:1424

- reduction of depressant effects of protamine sulfate on isolated ischemic rat hearts, 88:710
  - sensitivity of T-type calcium currents in various cell types, 88:168
  - vs sevoflurane, dynamic cerebral autoregulation, 88:341
  - vs sevoflurane, tracheal extubation, deeply anesthetized pediatric patients, 88:742
  - polyhalogenated methyl ethyl ethers, solubilities and anesthetic properties, 88:1161
- sevoflurane**
- anesthesia, followed by emergence "delirium," 88:1308
  - changes in biochemical markers of renal injury, healthy volunteers, 88:437
  - combined with N<sub>2</sub>O, subanesthetic concentrations, effect on analgesia, mood, and psychomotor performance, healthy volunteers, 88:1149
  - compound A, toxicity, disclosing of data used as basis for publication of studies (letter), 88:693
  - desflurane confusion and (letter), 88:1194
  - exposure, during four different methods of anesthetic induction, 88:925
  - in vitro* effects on platelet aggregation, 88:432
  - MAC, normal and cardiomyopathic hamsters, 88:489
  - N<sub>2</sub>O, mask induction, transient hyperdynamic response associated with controlled hypocapnic hyperventilation during, adults, 88:1384
  - N<sub>2</sub>O, trabecular outflow facility and formation rate of aqueous humor, rabbits, 88:781
  - recovery characteristics, preschool-aged children undergoing bilateral myringotomy and pressure equalization tube insertion, 88:34
  - remifentanyl, vs sevoflurane-N<sub>2</sub>O, trabecular outflow facility and formation rate of aqueous humor, rabbits, 88:781
  - renal effects, assessment using urinary markers, elderly patients (letter), 88:1426
  - subanesthetic concentrations, effect on analgesia, mood, and psychomotor performance, healthy volunteers, 88:1149

- vs isoflurane, dynamic cerebral autoregulation, 88:341
- vs isoflurane, tracheal extubation, deeply anesthetized pediatric patients, 88:742
- vs propofol, core hypothermia, 88:921
- vs propofol, effects on systemic toxicity of intravenous bupivacaine, rats, 88:155
- vs propofol, LMA insertion, adults, 88:908
- Anesthetic techniques**
  - acupressure, use to prevent PONV, meta-analysis, 88:1362
  - acupuncture, use to prevent PONV, meta-analysis, 88:1362
  - cerebrospinal drain, neurological complications, 88:388
  - combined spinal-epidural
    - epidural "top-up," effect of volume vs dose, 88:810
    - for labor analgesia, intrathecal sufentanil 10 µg plus bupivacaine 2.5 mg, effectiveness of half dose, 88:362
    - needle-through-needle technique, risk of metal toxicity, 88:393
  - continuous infusion, insulin, and neutrophil function, diabetic cardiac surgery patients, 88:1011
  - dorsal column stimulation, nationwide survey of academic teaching programs, 88:1311
  - electrical stimulation, nationwide survey of academic teaching programs, 88:1311
  - endobronchial insufflation catheter, for endobronchial laser surgery, 88:298
  - endoscopic thoracic sympathectomy, for primary erythromelalgia in upper extremities, 88:865
  - epidural
    - analgesia, and cesarean delivery, relationship (letter), 88:963
    - anesthetic management of familial hypokalemic periodic paralysis during parturition, 88:1081
    - catheter insertion, effects of needle bevel orientation, laboring parturients, 88:352
    - cesarean section, accidental dural puncture, cause of pneumocephalus with intense headache and unilateral pupillary dilatation, 88:582
    - cesarean section, and acid aspiration prophylaxis, German survey, 88:63
    - cesarean section, patient with Hippel-Lindau disease and multiple sclerosis, 88:1083
    - lidocaine, alkalinized, -induced hypotension, effect of epidural phenylephrine, 88:1322
    - lumbar, intensity of sensory block, effect of lidocaine concentration, 88:123
    - needle bevel, influence on spread of sensory blockade, laboring parturient (letter), 88:962
    - repetitive peridural administration of metoclopramide, histopathological changes after, dogs, 88:100
    - ropivacaine for initiation of labor epidural analgesia, dose-finding study, 88:1340
    - sufentanil analgesia, postoperative, effect of intravenous ketoprofen, children, 88:1036
    - test dose, epinephrine and diagnostic accuracy of aspiration during labor epidural analgesia, 88:1073
    - test dose, in obstetric patients, usefulness (editorial), 88:971
  - fiberoptic bronchoscopy, oral, simple technique (letter), 88:695
  - general
    - for cesarean section, and acid aspiration prophylaxis, German survey, 88:63
    - changes in maternal middle cerebral artery blood flow velocity, women with severe preeclampsia, 88:357
    - combined with thoracic epidural, vs general, arterial oxygenation during one-lung ventilation, 88:494
    - inducing, methods, onset time, recovery duration and drug cost, 88:930
    - quality of recovery score, development and psychometric testing, adults, 88:83
    - surgery under, urinary aquaporin-2 excretion and urine osmolality, effects of intrinsic vasopressin, 88:181
    - vs combined, arterial oxygenation during one-lung ventilation, 88:494
  - inhaled anesthetic, without muscle relaxant, tracheal intubation, healthy pediatric patients, survey of technique utilization and perceptions of safety, 88:737
  - intracerebroventricular morphine, for cancer pain management, CSF and plasma concentrations of morphine, morphine-3-glucuronide, and morphine-6-glucuronide, 88:109
  - intraoperative autotransfusion, small children, feasibility study, *in vitro* investigation, 88:763
  - intraspinal opioid therapy, nationwide survey of academic teaching programs, 88:1311
  - intrathecal opioids, for acute postoperative pain, safety and efficacy, 88:599
  - intravenous regional
    - adding clonidine to lidocaine, for prevention of tourniquet pain, 88:1327
    - meperidine, dose-response study, 88:831
    - peripheral antihyperalgesic effect of morphine, 88:117
  - in vitro* fertilization, addition of fentanyl to 1.5% lidocaine, 88:523
  - mask induction, sevoflurane-N<sub>2</sub>O, transient hyperdynamic response associated with controlled hypocapnic hyperventilation during, adults, 88:1384
  - millimeter waves, suppression of pain sensation, 88:836
  - monitored anesthesia care
    - remifentanyl administration during, intermittent boluses vs continuous infusion, 88:518
    - sedation during, cost-effectiveness of methohexital vs propofol, 88:723
  - neuraxial, temperature monitoring and management (editorial), 88:243
  - paravertebral blockade, for modified radical mastectomy, pregnant patient, 88:1351
  - patient-controlled analgesia (PCA)
    - morphine-6-glucuronide, persistently high plasma levels despite decreased hourly PCA morphine use after single-dose diclofenac, 88:1137
    - prophylactic antiemetics, efficacy and adverse effects, quantitative systematic review, 88:1354
  - spinal morphine and post-cesarean section analgesia, 88:346
  - percutaneous electrical nerve stimulation, analgesic response, effect of stimulus frequency, patients with chronic low back pain, 88:841
  - preemptive analgesia
    - anesthetics and spinal Fos, 88:414
    - preincisional dextromethorphan treatment, effect on postoperative

**Anesthetic techniques—continued**

- pain and opioid requirement after laparoscopic cholecystectomy, 88:1331
- regional
  - axillary block, clonidine, and postoperative pain, 88:1109
  - axillary block, repeated, and risk of persistent paresthesia, 88:382
  - axillary block, reversal of prolonged unconsciousness by naloxone after intravascular injection of local anesthetic, 88:1185
  - axillary brachial plexus blockade, duration, effect of tramadol added to mepivacaine, 88:853
  - brachial plexus block, new approach, under fluoroscopic guidance, 88:91
  - ilioinguinal nerve block, colonic puncture during, child, 88:1051
  - infraclavicular brachial plexus block, described by Raj et al., needle placement, lack of precision demonstrated by MRI, 88:593
  - interscalene block, prevention of autonomic hyperreflexia, quadriplegic patient undergoing shoulder surgery, 88:98
  - interscalene block, shoulder surgery, sitting position,  $\beta$ -adrenergic blockers and vasovagal episodes, 88:378
  - interscalene brachial plexus block, with 0.5% and 0.75% ropivacaine, pulmonary function changes, double-blinded comparison with 2% mepivacaine, 88:587
  - midfemoral block, new lateral approach to sciatic nerve (letter), 88:1426
  - neurological complications from CSF drains, 88:388
  - peripheral nerve block, nerve stimulator and multiple injection technique, failure rate, patient acceptance, and neurologic complications, 88:847
  - remifentanyl as adjunct sedative, safety and effectiveness, 88:134
  - supraclavicular nerve block, indirect vs direct measurement of brachial plexus, 88:1113
  - temperature monitoring practices during, 88:373
- spinal
  - analgesia, morphine and monoamine antinociception, rats, 88:141
  - lumbar dura mater biomechanics, experimental characterization and

- scanning electron microscopy observations, 88:1317
- meperidine, effects of three graded doses, African men, 88:827
- neurotoxicity of spinal drugs, 88:797
- patient with postdural puncture headache and acute leukemia, clinical dilemma, 88:166
- sameridine, safety and efficacy, comparative dose-ranging study with lidocaine for inguinal hernia repair, 88:815
- subarachnoid, assessing level of sensory block using pressure palpator, 88:398
- tetracaine and spinal cord ischemia, 88:56
- spinal cord stimulation, nationwide survey of academic teaching programs, 88:1311
- sympathectomy, and hyperbaric oxygen exposure, synergistic effect on transcutaneous  $PO_2$ , healthy volunteers, 88:67
- target-controlled infusion (TCI), -based anesthesia, vs standard anesthetic regimens, cost analysis, 88:77
- thoracic epidural
  - combined with general, vs general, arterial oxygenation during one-lung ventilation, 88:494
  - effects on intraoperative visceral perfusion and metabolism, 88:402
  - vs clonidine, perioperative stress response and release of troponin T, patients undergoing CABG, 88:701
- topical
  - EMLA, postburn, effect, digital image analysis of erythema development after experimental thermal injury to human skin, 88:1131
  - thoracostomy tube removal, 88:1107
- total intravenous anesthesia, TCI-based, vs standard anesthetic regimens, cost analysis, 88:77
- Angiotensin-converting enzyme inhibitors**, see Pharmacology
- Antagonists, neuromuscular relaxants**, neostigmine, neurotoxicity of spinal drugs, 88:797
- Antagonists, opioid**, naloxone, reversal of prolonged unconsciousness after intravascular injection of local anesthetic and clonidine, 88:1185
- Anterior cruciate ligament surgery**, see Surgery, knee
- Antibiotics**, isepamicin, chronic treatment with, resistance to mivacurium and rocuronium, rabbits, 88:654

**Antiemetics**

- preoperative oral, for reducing PONV after tonsillectomy, children, 88:1298
- prophylactic, during PCA, efficacy and adverse effects, quantitative systematic review, 88:1354
- Antinociception**, see Pain
- Aortic surgery**, see Surgery
- Aprotinin**, see Pharmacology
- Aquaporin-2 excretion**, see Urine
- Aqueous humor**, see Eye
- Arterial-enttidal carbon dioxide difference**, see Ventilation
- Arterialization**, dorsum vein on hand (letter), 88:1425
- Arterial thrombosis**, see Complications
- Arteries**
  - continuous intraarterial blood gas monitoring, reliability during one-lung ventilation (letter), 88:1192
  - gastroepiploic, human, contractile responses, endotoxin-induced attenuation, effects of vasopressin, 88:542
  - middle cerebral, blood flow velocity, maternal, changes associated with general anesthesia, women with severe preeclampsia, 88:357
  - oxygenation, during one-lung ventilation, combined vs general anesthesia, 88:494
  - retinal, air embolism, fluorescein angiographic evaluation of effects of hypotension and hemodilution, dogs, 88:1004
- Arterio-alveolar oxygen tension difference**, see Ventilation
- Arthroscopic meniscectomy**, see Surgery, knee
- ASC**, see Ambulatory surgery centers
- Aspiration**, see Complications
- Atracurium**, see Neuromuscular relaxants
- Atrial natriuretic peptide**, see Hormones
- Atrial pacing**, see Heart
- Audiometry**, see Measurement techniques
- Autonomic nervous system**,
  - cardiovascular autonomic dysfunction and hemodynamic response to anesthetic induction, patients with coronary artery disease and diabetes mellitus, 88:985
- Autotransfusion**, see Blood, transfusion
- Axillary block**, see Anesthetic techniques, regional
- Back surgery**, see Surgery
- Bacteria**, growth rates, in propofol and mixtures of propofol and lidocaine, 88:209
- Bacterial contamination**, see Infection



**Bain system**, see Equipment, anesthetic delivery systems

**Barbiturates**, see Theories of anesthetic action

**Benzocaine**, see Anesthetics, local

**Bispectral Index**, see Monitoring, electroencephalography

**Bleeding**, see Complications

**$\beta$ -Blockers**, see Pharmacology

## Blood

blood-brain barrier, osmotic, disruption, influence of anesthetic choice,  $\text{Paco}_2$ , and other factors, rats with brain tumor xenografts, 88:559

fibrinolysis, excess, during cardiac and noncardiac thoracic surgery, detection, use of latex agglutination D-dimer assay, 88:1226

hemodilution, effects on duration of retinal artery air embolism, dogs, 88:1004

## plasma

adrenomedullin concentration, increased, in hemorrhagic shock, dogs, 88:326

concentrations of morphine, morphine-3-glucuronide, and morphine-6-glucuronide, before and after initiation of intracerebroventricular morphine for cancer pain management, 88:109

## transfusions

autotransfusion, intraoperative, small children, feasibility, *in vitro* investigation, 88:763

requirements, in orthotopic liver transplantation (letter), 88:692  
requirements, reduction, effectiveness of thromboelastography-guided transfusion algorithm, complex cardiac surgery, 88:312

**Blood flow**, see Brain; Kidney; Spinal cord

## Blood pressure

effects of dexmedetomidine on neuromuscular blockade, human volunteers, 88:633

## hypertension

endothelin and major surgery, 88:898  
essential, patients with, recovering from surgery, effects of clonidine on sensitivity to phenylephrine and nitroprusside, 88:1239  
remifentanyl vs fentanyl and anesthetic induction, 88:734

## hypotension

effects on duration of retinal artery air embolism, dogs, 88:1004  
refractory, intraoperative, treatment with terlipressin, patients

chronically treated with antagonist of renin-angiotensin system, 88:980

increased, associated with controlled hypocapneic hyperventilation during sevoflurane- $\text{N}_2\text{O}$  mask induction, adults

maintained despite profound myocardial depression during acute bupivacaine overdose, pigs, 88:1143  
stress response to insertion of Combitube, LMA, or tracheal intubation, 88:1389

## Book and multimedia reviews

Anesthesia: Biologic Foundations, 88:235

Anesthesiology Pocket Guide, 88:236

Atlas of Anesthesia. Vol. II. Scientific Principles of Anesthesia., 88:471

The Cancer Patient. Vol. 16, No. 3 of Anesthesia Clinics of North America, 88:699

Churchill's Pocketbook of Pain Management, 88:968

Clinical Anesthesia Procedures of the Massachusetts General Hospital, 88:236

Drugs, Diseases, and Anesthesia, 88:472

The High Risk Obstetrical Patient. Vol. 16, No. 2 of Anesthesiology Clinics of North America, 88:471

The Laryngeal Mask Airway. Vol. 36, No. 2 of International Anesthesiology Clinics, 88:697

Management of Acute and Chronic Pain, 88:967

Management of the Operating Room. Vol. 36, No. 1 of International Anesthesiology Clinics, 88:699

The MGH Board Review of Anesthesiology: 5th ed., 88:1195

Pain Management. Vol. 6 in Atlas of Anesthesia, 88:159

Peripheral Nerve Blocks, Upper and Lower Limb, 88:697

Port-AccessT Cardiac Surgery for the Anesthesiologist, 88:235

Principles and Practice of Intensive Care Monitoring, 88:967

Principles of Anesthetic Techniques and Anesthetic Emergencies. Atlas of Anesthesia Series. Vol 4, 88:968

Simulators in Anesthesiology Education, 88:472

Total Intravenous Anesthesia, 88:698

**Brachial plexus**, see Anatomy; Nerve

**Brachial plexus block**, see Anesthetic techniques, regional

**Brachial plexus depth**, see Measurement techniques

## Brain

adrenomedullin, production after hypothermic CPB, adult cardiac surgical patients, 88:1030

blood-brain barrier, osmotic, disruption, influence of anesthetic choice,  $\text{Paco}_2$ , and other factors, rats with brain tumor xenografts, 88:559

## blood flow

oxygen delivery during retrograde cerebral perfusion, humans, 88:8  
sevoflurane vs isoflurane anesthesia, 88:341

velocity, during CPB, effects of propofol (letter), 88:231

velocity, middle cerebral artery, maternal, changes associated with general anesthesia, severe preeclampsia, 88:357

cortex, dexmedetomidine and anesthetic depth, 88: 625

dynamic autoregulation, sevoflurane vs isoflurane anesthesia, 88:341

## glutamate release

depolarization-evoked, different effects of volatile anesthetics and polyhalogenated alkanes, rat cortical brain slices  
tetracaine and spinal cord ischemia, rabbits, 88:56

## hippocampus

area CA1, excitatory and inhibitory synaptic transmission, effects of general anesthetics, 88:676  
microdialysis, anesthetics, catecholamines, and cerebral ischemia, 88:787

hypothalamus, dexmedetomidine and CNS pressor responses, 88:617

intracranial pressure, and venous cannulation, CPB, 88:329

## ischemia

focal, brain injury after, and fentanyl, rats, 88:49

near-complete, concentrations of brain and plasma catecholamines, effects of isoflurane, ketamine, and fentanyl/ $\text{N}_2\text{O}$ , rats, 88:787

outcome, effects of aprotinin, rats, 88:1  
oxygen delivery during retrograde cerebral perfusion, humans, 88:8

metabolism, oxygen delivery during retrograde cerebral perfusion, humans, 88:8

motoneurons, excitability, effect of isoflurane, F-wave study

**Brain—continued**

- oxygenation
  - continuous measurement by near infrared spectroscopy during induction of anesthesia, 88:554
  - monitoring, after traumatic brain injury, measuring brain tissue oxygenation vs jugular venous oxygen saturation, 88:549
- oxygen consumption, oxygen delivery during retrograde cerebral perfusion, humans, 88:8
- oxygen extraction, during CPB, effects of propofol (letter), 88:231
- raphe magnus, activity of neurons, effect of  $\alpha$ -adrenergic drugs, rats, 88:459
- reticular formation, dexmedetomidine and CNS pressor responses, 88:617
- rostral ventrolateral medulla, activation during strychnine allodynia, 88:1125
- tumor, influence of anesthetic choice,  $\text{Paco}_2$ , and other factors on osmotic blood-brain barrier disruption, 88:559
- water, saline vs lactated Ringer's, human volunteers, 88:999

**Brain injury, see Complications****Breath sounds, see Examination****Bronchial rupture, see Complications****Bronchoscopic swivel adapter, see Equipment****Bupivacaine, see Anesthetics, local****Burns, see Complications****CABG, see Surgery, coronary artery bypass grafting****Calcium, see Ions****Cancer, see Pain****Capnography, see Monitoring****Carbon dioxide**

- hypoxic ventilatory response, effects of  $\text{CO}_2$  and of sustained hypoxia (letter), 88:695

**insufflation**

- peritoneal, pneumoperitoneum, hemodynamic effects, assessment by continuous esophageal aortic blood flow echo Doppler, healthy infants (letter), 88:468

- retroperitoneal, cardiopulmonary dysfunction during minimally invasive thoraco-lumboendoscopic spine surgery, 88:4688:1244

- $\text{Paco}_2$ , influence on osmotic blood-brain barrier disruption, rats with brain tumor xenografts, 88:559

- tension, transcutaneous monitoring after cardiothoracic surgery, infants and children, 88:531

**Carbon dioxide embolism, see Complications****Cardiac arrest, see Complications****Cardiomyopathy, see Heart****Cardiopulmonary bypass, see Surgery****Cardiopulmonary dysfunction, see Complications****Cardiothoracic surgery, see Surgery****Catecholamines, see Sympathetic nervous system****Catheters, see Equipment****Cells, neutrophils****function**

- effect of continuous insulin infusion, diabetic cardiac surgery patients, 88:1011

- effects of clonidine and dexmedetomidine, humans, 88:452

- postischemic adherence, coronary system of isolated guinea pig hearts, effects of S(+)-ketamine vs R(-)-ketamine, 88:1017

**Central nervous system (CNS), pressor sites, stimulation, cardiovascular responses, modulation by dexmedetomidine, 88:617****Cerebrospinal drain, see Anesthetic techniques****Cerebrospinal fluid (CSF)**

- concentrations of morphine, morphine-3-glucuronide, and morphine-6-glucuronide, before and after initiation of intracerebroventricular morphine for cancer pain management, 88:109

- drainage, neurological complications, 88:388

**Cesarean section, see Surgery****Cholecystectomy, see Surgery****Cisatracurium, see Neuromuscular relaxants****Clonidine, see Analgesics; Pharmacology; Receptors,  $\alpha_2$ -adrenergic, agonists****Closed head trauma, see Complications****Cloth band, see Complications****Coagulation**

- heparin, direct effects on canine smooth muscle tone, dogs, 88:1232

- markers, effects of *in vitro* hemodilution with gelatin, hydroxyethyl starch, and lactated Ringer's solution, analysis using SONOCLOTT, 88:483

**platelets**

- aggregation, *in vitro* effects of isoflurane, sevoflurane, and propofol, 88:432

- function, after cardiac surgery, effect of intravenously administered magnesium, 88:1213

- hyporeactivity, during CPB, young infants, 88:258

- protamine, direct effects on canine smooth muscle tone, dogs, 88:1232

**Cold pressor test, see Measurement techniques****Colloids**

- gelatin, effects of *in vitro* hemodilution on coagulation markers, analysis using SONOCLOTT, 88:483

- Hextend, randomized phase III clinical trial, 88:992

- hydroxyethyl starch, effects of *in vitro* hemodilution on coagulation markers, analysis using SONOCLOTT, 88:483

**Colonial puncture, see Complications****Combined spinal-epidural, see Anesthetic techniques****Combitube, see Equipment****Common peroneal nerve, see Nerve****Complications**

- acid aspiration syndrome, prophylaxis, and anesthesia for cesarean section, German survey, 88:63

- acute myocardial infarction, during lung volume reduction surgery, 88:332

- after ambulatory surgery, 88:508

- alcohol withdrawal syndrome, in surgical patients, prevention and treatment, 88:946

- amniotic fluid embolism, and inhaled NO (letter), 88:691

- anaphylactic reaction, severe, to cisatracurium, 88:462

- arterial thrombosis, acute postoperative, protein S deficiency presenting as, four-year-old child, 88:535

- aspiration, liquid flow with LMA, 88:904

**bleeding**

- hemorrhagic shock, increased plasma concentration of adrenomedullin, dogs, 88:326

- tracheo-innominate artery fistula after tracheostomy, 88:777

- brain injury, traumatic, monitoring cerebral oxygenation after, measuring brain tissue oxygenation vs jugular venous oxygen saturation, 88:549

- bronchial rupture, by double-lumen endobronchial tube, during staging thoracoscopy, 88:1252

- burns, postburn topical EMLA, effect, digital image analysis of erythema development after experimental thermal injury to human skin, 88:1131

- carbon dioxide embolism, infant (letter), 88:227

- cardiac arrest, tracheo-innominate artery fistula after tracheostomy, 88:777
- cardiopulmonary dysfunction, during minimally invasive thoracolumboendoscopic spine surgery, 88:1244
- closed head trauma, use of vasopressors after (letter), 88:1423
- cloth band, as cause of difficult endotracheal intubation (letter), 88:230
- colonic puncture, during ilioinguinal nerve block, child, 88:1051
- coronary artery disease, patients with atrial natriuretic peptide, attenuation of pacing-induced myocardial ischemia during general anesthesia, 88:279
- cardiovascular autonomic dysfunction and hemodynamic response to anesthetic induction, 88:985
- effect of nitroglycerin on pacing-induced changes in myocardial oxygen consumption and metabolic coronary vasodilation, 88:271
- diabetes
  - cardiac surgery patients, continuous insulin infusion, effect on neutrophil function, 88:1011
  - coronary artery disease and, patients with, cardiovascular autonomic dysfunction and hemodynamic response to anesthetic induction, 88:985
- double-lumen endotracheal tube obstruction, unusual case (letter), 88:694
- emergence "delirium," after sevoflurane anesthesia, 88:1308
- emphysema, lung volume reduction surgery, preoperative functional predictors for postoperative outcome, 88:28
- erythromelalgia, primary, in upper extremities, endoscopic thoracic sympathectomy, 88:865
- failed back surgery syndrome
  - treatment, use of epidural steroids, cost-benefit of fluoroscopy, 88:467
  - treatment, use of epidural steroids (editorial), 88:240
- familial hypokalemic periodic paralysis, anesthetic management during parturition, 88:1081
- fat embolism, and neurological dysfunction (letter), 88:1427
- gas embolism, during laparoscopic nephrectomy, detection by TEE, 88:500
- headache
  - intense, after accidental dural puncture during epidural anesthesia for cesarean section, 88:582
  - postdural puncture, patient with acute leukemia, clinical dilemma, 88:166
- hematomas, epidural, associated with enoxaparin (letter), 88:229
- hyperreflexia, autonomic, prevention, use of interscalene regional anesthesia, 88:98
- hypomagnesia, after cardiac surgery, effect of intravenously administered magnesium, 88:1213
- hypotension, induced by alkalinized epidural anesthesia, effect of epidural phenylephrine, 88:1322
- hypoxemia, early postoperative, influence of surgical sites, adults undergoing elective surgery, 88:213
- intraperitoneal adhesions, effect of tenoxicam, mice, 88:939
- intravascular injection, detecting, T-wave amplitude as guide, anesthetized children, 88:754
- leukemia, acute, patient with postdural puncture headache, clinical dilemma, 88:166
- macroglossia, postoperative, rapid resolution after extubation of trachea, 88:220
- malignant hyperthermia susceptibility, in humans, diagnosis, European vs North American protocols, 88:1155
- metal toxicity, risk, in combined spinal-epidural anesthesia, 88:393
- methemoglobinemia, severe, suspected, caused by topical application of ointment containing benzocaine around enterostomy (letter), 88:1190
- nerve injury, after placement of CSF drainage catheters and needles, anesthetized patients, implications for regional anesthesia, 88:388
- neurological
  - and fat embolism (letter), 88:1427
  - nerve stimulator for peripheral regional anesthesia, 88:847
  - osteopetrosis, malignant, patients with, risk of anesthetic morbidity and mortality, 88:1292
  - panic disorder, and lactated Ringer's solution, relationship, 88:795
  - paresthesia, persistent, risk, and repeated axillary block, 88:382
  - pheochromocytoma, laparoscopic adrenalectomy for, hemodynamic changes and catecholamine release during, 88:16
- pneumocephalus, with intense headache and unilateral pupillary dilation, after accidental dural puncture during epidural anesthesia for cesarean section, 88:582
- pneumothorax, risk, associated with supraclavicular nerve block, indirect vs direct measurement of brachial plexus depth, 88:1113
- preeclampsia, severe, women with, changes in maternal middle cerebral artery blood flow velocity, 88:357
- prolonged unconsciousness, after intravascular injection of local anesthetic and clonidine, reversal by naloxone, 88:1185
- protein S deficiency, presenting as acute postoperative thrombosis, four-year-old child, 88:535
- pulmonary embolism
  - hemodynamic changes after, effects of small-dose inhaled NO, dogs, 88:1025
  - massive intraoperative, and succinylcholine (letter), 88:1423
  - massive, nitric oxide inhalation during (letter), 88:1188
- pulmonary hypertension
  - during protamine infusion after CPB, treatment, use of NO, 88:505
  - small-dose inhaled NO and hemodynamic changes after pulmonary air embolism, dogs, 88:1025
- reflex sympathetic dystrophy, rat paw perfusion after mononeuropathy, 88:103
- regurgitation, liquid flow with LMA, 88:904
- respiratory arrests, and epidural opioids (letter), 88:962
- shivering
  - postanesthetic, treatment of, comparison among nalbuphine, meperidine, and placebo, 88:686
  - postoperative, effects of prostaglandin E<sub>1</sub>, neurosurgical patients undergoing deliberate mild hypothermia, 88:446
- shock, hemorrhagic, increased plasma concentration of adrenomedullin, dogs, 88:326
- systemic inflammatory response syndrome, due to intraperitoneal hyperthermia, lung injury associated with, prevention by methylprednisolone, 88:771

**Complications—continued**

- tachycardia
    - induced subendocardial necrosis, acutely instrumented dogs with fixed coronary stenosis, 88:973
    - pacemaker-driven (letter), 88:965
    - perioperative torsade de pointes ventricular (letter), 88:966
  - tracheo-innominate artery fistula, after tracheostomy, 88:777
  - upper respiratory tract infections, children with, use of LMA (letter), 88:693
  - vasovagal episodes, and  $\beta$ -adrenergic blockers, during shoulder surgery in sitting position during interscalene block, 88:378
  - venous air embolism, during Parkinson surgery, patients with spontaneous ventilation, 88:793
  - von Hippel-Lindau disease, and multiple sclerosis, patient with, epidural anesthesia for cesarean section, 88:1083
- Composite sedation**, see Measurement techniques
- Compound A**, see Anesthetics, volatile, sevoflurane; Metabolism
- Computed tomographic scans**, see Monitoring
- Computer-assisted auscultation**, see Measurement techniques
- Computer-assisted image analysis**, see Measurement techniques
- Computer simulation**
- to determine effect of rapid anesthetic recovery protocols on ASC staffing, 88:1053
  - study, operating room efficiency, decrease in case duration required to complete an additional case, 88:72
- Continuous esophageal aortic blood flow echo Doppler**, see Monitoring
- Convulsants, gaseous**, polyhalogenated, convulsant inhaled compounds, 88:884
- Convulsants, volatile**, polyhalogenated, convulsant inhaled compounds, 88:884
- Coronary artery bypass grafting**, see Surgery
- Coronary artery disease**, see Complications
- Cortex**, see Brain
- Cost**, see Economics
- CPB**, see Surgery, cardiopulmonary bypass
- Cranial trauma**, see Complications, closed head trauma
- Craniotomy procedures**, see Surgery

**Critical care and trauma**, see Anesthesia**Critical care medicine**

- alcohol withdrawal syndrome, in surgical patients, prevention and treatment, 88:946
  - journals, important, national origin of articles, MEDLINE-based analysis, 88:1175
- Crystalloids**, Ringer's lactate, effects of *in vitro* hemodilution on coagulation markers, analysis using SONOCLOTT, 88:483
- CSF**, see Cerebrospinal fluid
- Cuffed oropharyngeal airway**, see Equipment

**Data**, used as basis for publication, disclosing of (letter), 88:693**D-dimer assay**, see Measurement techniques**Descending inhibitory system**, see Spinal cord**Desflurane**, see Anesthetics, volatile**Desipramine**, see Pharmacology**Dexamethasone**, see Antiemetics**Dexmedetomidine**, see Anesthetics, intravenous; Pharmacology**Dextromethorphan**, see Pharmacology; Premedication**Diabetes**, see Complications**1,2-Dichlorohexafluorocyclobutane**, see Polyhalogenated alkanes**Diclofenac**, see Pharmacology**Discharge**

- criteria, after ambulatory surgery, 88:508
- early, after laparoscopic cholecystectomy, patient preferences, 88:1280

**Dorsal column stimulation**, see Anesthetic techniques**Dorsum veia**, see Veins**Double-lumen tubes**, see Equipment**Droperidol**, see Pharmacology**Dura mater**, lumbar, biomechanics, experimental characterization and scanning electron microscopy observations, 88:1317**Dynamic compliance**, see Lung**Ear**, audiological reaction times, effects of midazolam, human volunteers, 88:1064**Economics**

- cost
  - drug, comparison of induction methods, 88:930
  - due to waiting time, intense neuromuscular block for anesthesia during endolaryngeal procedures, 88:1335

- effectiveness, methohexital vs propofol for sedation during monitored anesthesia care, 88:723
- cost analysis, target-controlled infusion, vs standard anesthetic regimens, 88:77
- cost-effectiveness, ratio (letter), 88:1191
- health systems research and financial impact of decreasing cost duration, 88:72
- quality of recovery score, 88:83
- target-controlled infusion and costs, 88:77
- operating room efficiency, decrease in case duration required to complete an additional case, computer simulation study, 88:72

**Education**

- fellowship, academic teaching programs, nationwide survey, state of implantable pain therapies, United States, 88:1311

**patient**

- parental education and anxiolysis, facilitation, use of preanesthetic video before pediatric ambulatory surgery, 88:246
- parents' preoperative anxiety and anesthesia knowledge, measures, validation, 88:251

**residents**

- clinical performance, relationship of cognitive, personality and academic measures, 88:1092
- valuing work performed by (letter), 88:964

- training courses, Interventional Training Workshop, for implantable pain therapies (editorial), 88:1203

**EE**, see Monitoring, epicardial**echocardiography****EEG**, see Monitoring, electroencephalography**Electrical stimulation**, see Anesthetic techniques**Electrocardiography**, see Monitoring**Electrocorticography**, see Monitoring**Electroencephalography**, see Monitoring**Emergence "delirium,"** see Complications**Emergency medicine**, journals, important, national origin of articles, MEDLINE-based analysis, 88:1175**EMLA cream**, see Anesthetics, local**Emphysema**, see Complications**Endobronchial insufflation catheter**, see Anesthetic techniques**Endobronchial tube**, see Equipment**Endolaryngeal procedures**, see Surgery**Endoscopic thoracic sympathectomy**, see Anesthetic techniques**Endothelin-1**, see Pharmacology

- Endotoxins**, see Toxins
- Enflurane**, see Anesthetics, volatile
- Enoxaparin**, see Pharmacology
- Epidural anesthesia**, see Anesthetic techniques; Anesthetic techniques, thoracic epidural
- Epilepsy**, see Surgery
- Epinephrine**, see Pharmacology; Sympathetic nervous system, catecholamines
- Equipment**
- anesthesia machines, technique for filling vaporizer (letter), 88:233
  - anesthetic delivery systems, Bain system, vs adult circle breathing system, infant lung model, 88:527
  - bronchoscopic swivel adapter, another use (letter), 88:1187
  - catheters
    - CSF drainage, placement, neurological complications after, 88:388
    - double-lumen jet, respiratory gas monitoring during high-frequency jet ventilation for tracheal resection, 88:224
    - epidural, insertion, effect of needle bevel orientation, laboring parturients, 88:352
  - Combitube
    - as emergency airway device for anesthesiologists (letter), 88:233
    - esophageal-tracheal, fiberoptic-guided airway exchange, spontaneously breathing vs mechanically ventilated patients, 88:193
    - insertion, hemodynamic and catecholamine stress responses, 88:1389
  - cuffed oropharyngeal airway
    - repeated use in infant for radiation therapy (letter), 88:1421
    - vs LMA, alternative hypotheses (letter), 88:961
    - vs LMA, target plasma concentration of propofol required to place, 88:913
  - double-lumen endobronchial tube
    - bronchial rupture during staging thoracoscopy, 88:1252
    - size, predicting, use of computed tomographic scan measurements of left main bronchus diameter, 88:302
  - double-lumen endotracheal tube, obstruction, unusual case (letter), 88:694
  - double-lumen jet catheter, respiratory gas monitoring during high-frequency jet ventilation for tracheal resection, 88:224
  - double-lumen tubes, selecting, for small patients (letter), 88:466
  - endotracheal tube, Murphy eye, and reliability of chest auscultation in detecting endobronchial intubation, 88:1380
  - esophageal detector device, modified (letter), 88:691
  - fiberoptic bronchoscope, stuck, saline solution as lubrication (letter), 88:965
  - glottis simulator (letter), 88:1422
  - Guedel airway, modified, adjunct to unanticipated difficult airway (letter), 88:468
  - illuminating stylet, as aid for tracheal intubation via intubating LMA (letter), 88:470
  - laryngeal mask airway (LMA)
    - disposable, vs reusable, mucosal pressure, mechanism of seal, airway sealing pressure, and anatomic position, 88:1418
    - easy fiberoptic intubation, concept (letter), 88:228
    - effect on liquid flow between esophagus and pharynx, randomized, controlled cadaver study, 88:904
    - flexible and standard, oropharyngeal leak pressure and cuff position, influence of head and neck position, 88:913
    - flexible reinforced LMA, vs tracheal intubation, anesthesia for intranasal surgery, 88:421
    - insertion, hemodynamic and catecholamine stress responses, 88:1389
    - insertion, sevoflurane vs propofol, adults, 88:908
    - intubating, tracheal intubation, illuminating stylet as aid (letter), 88:470
    - intubating, ventilation difficulties, 88:1181
    - intubating, with and without fiberoptic guidance, 88:662
    - reusable, vs disposable, mucosal pressure, mechanism of seal, airway sealing pressure, and anatomic position, 88:1418
    - undetected hole (letter), 88:232
    - use before tracheal intubation, patient with cancerous nose, 88:1429
    - use during emergence from anesthesia, patient with unstable neck (letter), 88:469
    - use in children with upper respiratory tract infections (letter), 88:693
    - vs cuffed oropharyngeal airway, alternative hypotheses (letter), 88:961
    - vs cuffed oropharyngeal airway, target plasma concentration of propofol required to place, 88:913
  - laryngoscope, fractured bulb (letter), 88:1424
  - laryngotracheal topical anesthesia kit, with capnography, for difficult endotracheal intubation, 88:943
  - Macintosh blade, modified, difficult airway management with balloon inflation (letter), 88:1425
  - needles
    - bevel orientation, effects during epidural catheter insertion, laboring parturients, 88:352
    - CSF drainage, placement, neurological complications after, 88:388
    - epidural, bevel, influence on spread of sensory blockade, laboring parturient (letter), 88:962
    - MRI and Raj's infraclavicular block, 88:587
  - pacemakers, and tachycardia (letter), 88:965
  - pressure palpator, assessment of level of sensory block after subarachnoid anesthesia, 88:398
  - pulse oximeter, false comfort from (letter), 88:1428
  - vaporizers
    - desflurane, use of electricity (letter), 88:1189
    - level indicator, obstruction, detected by anesthetic vapor analysis (letter), 88:469
  - ventilators
    - Ohmeda 7900, flow sensor, protecting (letter), 88:234
    - Ohmeda 7900, vs Ohmeda 7000, changes in fresh gas flow, effects on volume-controlled ventilation, 88:39
- Erythromelalgia**, see Complications
- Esophageal detector device**, see Equipment
- Esophageal monitoring**, see Monitoring
- Etomidate**, see Anesthetics, intravenous
- European Malignant Hyperthermia Group**, protocols for diagnosis of malignant hyperthermia susceptibility, humans, 88:1155
- Examination**, breath sounds, reliability in detecting endobronchial intubation, and Murphy eye, 88:1380



- Extubation**, tracheal, of deeply anesthetized pediatric patients, isoflurane vs sevoflurane, 88:742
- Eye**
- aqueous humor, formation rate, sevoflurane-N<sub>2</sub>O vs sevoflurane-remifentanyl, rabbits, 88:781
  - intraocular compliance, sevoflurane-N<sub>2</sub>O vs sevoflurane-remifentanyl, rabbits, 88:781
  - pupillary dilatation, unilateral, after accidental dural puncture during epidural anesthesia for cesarean section, 88:582
  - retinal artery air embolism, fluorescein angiographic evaluation of effects of hypotension and hemodilution, dogs, 88:1004
- Failed back surgery syndrome**, see Complications
- Familial hypokalemic periodic paralysis**, see Complications
- Fast-tracking**, see Recovery
- Fat embolism**, see Complications
- Fellowship**, see Education
- Fentanyl**, see Analgesics; Anesthetics, intravenous
- Fiberoptic bronchoscope**, see Equipment
- Fiberoptic bronchoscopy**, see Anesthetic techniques
- Fibrinolysis**, see Blood
- Fluid balance**
- hypovolemia, Hextend, randomized phase III clinical trial, 88:992
  - lactated Ringer's solution and panic attacks, relationship, 88:795 vs 0.9% sodium chloride solution, effect on serum osmolality, human volunteers, 88:999
  - sodium chloride, 0.9% solution, vs lactated Ringer's solution, effect on serum osmolality, human volunteers, 88:999
- Fluorescein angiography**, evaluation of effects of hypotension and hemodilution on duration of retinal artery air embolism, dogs, 88:1004
- Fluorinated alkanols**, see Anesthetics, intravenous; Anesthetics, volatile
- Fluoroscopy**, and epidural steroids, failed back surgery syndrome, 88:367
- Formalin test**, see Anesthetic techniques
- F-wave study**, effect of isoflurane on motoneuron excitability, 88:681
- GABA receptors**, see Receptors
- Gas embolism**, see Complications
- Gases, nonanesthetic**, nitric oxide (NO) inhalation, during massive pulmonary embolism (letter), 88:1188
- inhaled, and amniotic fluid embolism (letter), 88:691
- management of catastrophic pulmonary vasoconstriction arising from protamine administration, 88:505
- small-dose inhaled, effects on hemodynamic changes after pulmonary air embolism, dogs, 88:1025
- Gastric tonometry**, see Measurement techniques
- Gastroepiploic arteries**, see Arteries
- Gastrointestinal system**, gastric emptying, inhibition by naloxone, rats, 88:204
- Gelatin**, see Colloids
- General anesthesia**, see Anesthetic techniques
- Glottis simulator**, see Equipment
- Glutamate release**, see Brain; Spinal cord, synaptosomes
- Granisetron**, see Antiemetics
- Guedel airway**, see Equipment
- Halothane**, see Anesthetics, volatile
- Hand**, dorsum vein, arterialization (letter), 88:1425
- H<sub>2</sub> blocking drugs**, see Pharmacology
- Head**, see Position
- Headache**, see Complications
- Heart**
- atrial pacing
    - induced changes in myocardial oxygen consumption and metabolic coronary vasodilation, patients with coronary artery disease, 88:271
    - induced myocardial ischemia during general anesthesia, effect of atrial natriuretic peptide, patients with coronary artery disease, 88:279
  - cardiomyopathy, MAC of volatile anesthetics, hamsters, 88:489
  - ischemia
    - cardioplegic, isoflurane, nitroprusside, and protamine interactions 88:710
    - effect of heart rate control among high-risk patients after vascular surgery, 88:47588:477
  - pacing-induced, during general anesthesia, attenuation by atrial natriuretic peptide, patients with coronary artery disease, 87:279
  - postischemic adherence of neutrophils, S(+)-ketamine vs R(-)-ketamine, guinea pigs
  - tachycardia-induced, subendocardial necrosis, acutely instrumented dogs with fixed coronary stenosis, 88:973
  - thoracic epidural anesthesia, clonidine, and CABG, 88:701
- mitral valve
- regurgitation, evaluation of location and mechanism, improved, with systematic TEE examination, 88:1205
  - systematic TEE examination, evolution into twenty-first century (editorial), 88:1197
- myocardial depression, profound, during acute bupivacaine overdose, pigs, 88:1143
- oxygen consumption, pacing-induced changes, effect of nitroglycerin, patients with coronary artery disease, 88:271
- rate**
- changes, fetal, incidence, after intrathecal fentanyl labor analgesia, 88:577
  - control, effects on myocardial ischemia among high-risk patients after vascular surgery, 88:47588:477
  - control, patient-based target-specific, individualizing  $\beta$ -adrenergic blocker therapy (editorial), 88:475
  - effects of dexmedetomidine on neuromuscular blockade, human volunteers, 88:633
  - tachycardia, associated with controlled hypocapnic hyperventilation during sevoflurane-N<sub>2</sub>O mask induction, adults
  - variability, during anesthetic induction, patients with coronary artery disease and diabetes mellitus, 88:985
- right-to-left shunt, effect on rate of rise of arterial and end-tidal halothane, children, 88:759
- sounds, amplitude and frequency characteristics, effect of halothane, children, 88:263
- tumor, resection, intraoperative management with aid of TEE and EE, pediatric patient, 88:1048
- valvular dysfunction, during laparoscopic nephrectomy, detection by TEE, 88:500
- vasodilation, pacing-induced changes, effect of nitroglycerin, patients with coronary artery disease, 88:271
- Hematomas**, see Complications
- Hemodilution**, see Blood
- Hemorrhagic shock**, see Complications, shock
- Heparin**, see Coagulation
- Hernia repair**, see Surgery
- Hextend**, see Colloids
- Hip surgery**, see Surgery

- Histopathology**, changes, after repetitive peridural administration of metoclopramide, dogs, 88:100
- Holter monitoring**, see Monitoring
- Hormones**, atrial natriuretic peptide, attenuation of pacing-induced myocardial ischemia during general anesthesia, patients with coronary artery disease, 88:279
- Hydroxyethyl starch**, see Colloids
- Hyperalgesia**, see Pain
- Hyperreflexia**, see Complications
- Hypertension**, see Blood pressure; Lung, pulmonary hypertension
- Hyperthermia**, malignant, susceptibility, in humans, diagnosis, European vs North American protocols, 88:1155
- Hypomagnesia**, see Complications
- Hypotension**, see Blood pressure; Complications
- Hypothalamus**, see Brain
- Hypothermia**  
core, sevoflurane vs propofol, 88:921  
deliberate mild, for neurosurgical procedures, effects of prostaglandin E<sub>1</sub> on intraoperative temperature changes and incidence of postoperative shivering, 88:446  
endothelin and major surgery, 88:898  
hypothermic circulatory arrest, brain metabolism, 88:8  
hypothermic CPB, cerebral production of adrenomedullin after, adult cardiac surgical patients, 88:1030  
moderate  
alveolar-arterial difference in oxygen tension increases with temperature-corrected determination, 88:538  
transient spinal cord ischemia and, rabbits, 88:56
- Hypovolemia**, see Fluid balance
- Hypoxemia**, see Complications
- Hypoxia**  
during minimally invasive thoracolumboendoscopic spine surgery, 88:1244  
endobronchial intubation as cause of immediate increase in peak inflation pressure, pediatric patients, 88:268  
hypoxic ventilatory response, effects of CO<sub>2</sub> and of sustained hypoxia (letter), 88:695
- Ilioinguinal nerve block**, see Anesthetic techniques, regional
- Illuminating stylet**, see Equipment
- Imaging techniques**, magnetic resonance imaging (MRI), demonstration of lack of precision in needle placement by infraclavicular brachial plexus block described by Raj et al., 88:593
- Immunology**, neutrophil functions, effects of clonidine and dexmedetomidine, humans, 88:452
- Implantable devices**, see Pain, therapies
- Induction**  
continuous measurement of cerebral oxygenation by near infrared spectroscopy, 88:554  
distress during, and postoperative behavioral outcomes, 88:1042  
hemodynamic response, patients with coronary artery disease and diabetes mellitus, 88:985  
methods, exposure to sevoflurane and N<sub>2</sub>O, 88:925  
remifentanyl vs fentanyl, ambulatory setting, 88:734  
technique  
and core temperature, 88:921  
onset time, recovery duration and drug cost, 88:930
- Infection**, bacterial contamination, of propofol, effect of lidocaine, 88:936
- Inflammation**, peripheral antihyperalgesic effects of morphine, healthy volunteers, after ultraviolet-B irradiation, 88:117
- Infraclavicular brachial plexus block**, see Anesthetic techniques, regional
- Injection**, see Pain
- Inspiratory flow**, see Lung
- Interactions, drug**  
isepamicin  
and mivacurium, 88:654  
and rocuronium, 88:654  
isoflurane, and protamine, 88:710  
mivacurium, and isepamicin, 88:654  
propofol, and thiopental, 88:667  
protamine  
and isoflurane, 88:710  
and sodium nitroprusside, 88:710  
rocuronium, and isepamicin, 88:654  
sodium nitroprusside, and protamine, 88:710  
thiopental, and propofol, 88:667
- Interscalene block**, see Anesthetic techniques, regional
- Interventional Training Workshop**, see Education, training courses
- Intracerebroventricular morphine**, see Anesthetic techniques
- Intranasal surgery**, see Surgery
- Intraocular compliance**, see Eye
- Intraperitoneal adhesions**, see Complications
- Intraspinal opioid therapy**, see Anesthetic techniques
- Intrathecal opioids**, see Anesthetic techniques
- Intravascular injection**, see Complications
- Intravenous regional anesthesia**, see Anesthetic techniques
- Intubation**  
endobronchial  
cause of immediate increase in peak inflation pressure, pediatric patients, 88:268  
detecting, reliability of chest auscultation, and Murphy eye, 88:1380  
infants and small children (letter), 88:229  
selecting size of double-lumen endobronchial tubes, 88:302  
endotracheal  
difficult, cloth band as cause (letter), 88:230  
difficult, laryngotracheal topical anesthesia kit with capnography, 88:943  
fiberoptic  
easy, via LMA, concept (letter), 88:228  
intubating LMA with and without fiberoptic guidance, 88:662  
nasotracheal, postoperative macroglossia, rapid resolution after extubation of trachea, 88:220  
remifentanyl vs fentanyl and anesthetic induction, 88:734  
tracheal  
esophageal-tracheal Combitube, fiberoptic-guided airway exchange, spontaneously breathing vs mechanically ventilated patients, 88:193  
healthy pediatric patients, without muscle relaxants, survey of technique utilization and perceptions of safety, 88:737  
hemodynamic and catecholamine stress responses, 88:1389  
technique, verifying tracheal tube placement, assessment of four different methods, critical care setting, 88:766  
use of LMA, patient with cancerous nose, 88:1429  
ventilation difficulties through intubating LMA, 88:1181  
via intubating LMA, illuminating stylet as aid (letter), 88:470  
vs flexible reinforced LMA, anesthesia for intranasal surgery, 88:421
- In vitro contracture test**, see Measurement techniques
- In vitro fertilization**, see Anesthetic techniques
- Ions**  
calcium

**Ions—continued**

- calcium-channel blockers, nifedipine, attenuation of hemodynamic aberrances during laparoscopic adrenalectomy for pheochromocytoma, 88:16
- heparin and protamine effects on tracheal smooth muscle, 88:1232
- T-type calcium currents, volatile anesthetic sensitivity in various cell types, 88:168
- magnesium, intravenously administered, effect on platelet function after cardiac surgery, 88:1213

**Ischemia**, see Brain; Heart; Spinal cord

**Isepalamicin**, see Antibiotics

**Isoflurane**, see Anesthetics, volatile

**Isoproterenol**, see Pharmacology

**Jet ventilation**, see Ventilation

**Jugular vein**, see Veins

**Jugular venous oxygen saturation**, see Monitoring

**Ketamine**, see Anesthetics, intravenous

**Ketoprofen**, see Analgesics

**Ketorolac**, see Analgesics

**Kidney**

- blood flow, diclofenac does not decrease, elderly patients undergoing orthopedic surgery, 88:149
- effects of sevoflurane, assessment using urinary markers, elderly patients (letter), 88:1426
- glomerular filtration, diclofenac does not decrease, elderly patients undergoing orthopedic surgery, 88:149
- renal injury, sevoflurane, disclosing of data used as basis for publication of studies (letter), 88:693
- toxicity, Compound A and renal injury, humans, 88:437

**Knee surgery**, see Surgery

**Labeling**, see Pharmacology

**Labor pain**, see Pain

**Lactated Ringer's solution**, see Crystalloids, Ringer's lactate; Fluid balance

**Laparoscopic surgery**, see Surgery

**Laryngeal mask airway**, see Equipment

**Laryngoscope**, see Equipment

**Laryngoscopy**, remifentanyl vs fentanyl and anesthetic induction, 88:734

**Laryngotracheal topical anesthesia kit**, see Equipment

**Larynx**, laryngeal adductors, assessing neuromuscular block, 88:426

**Laser Doppler fluxmetry**, see Measurement techniques

**Laser surgery**, see Surgery

**Leukemia**, see Complications

**Lidocaine**, see Anesthetics, local

**Liver transplantation**, see Surgery

**LMA**, see Equipment, laryngeal mask airway

**Low back pain**, see Pain

**Lung**

- dynamic compliance, and inspiratory flow, relationship (letter), 88:465
- injury, decreased tumor necrosis factor- $\alpha$  and inflammatory syndrome by steroids and intraperitoneal hyperthermia perfusion, 88:771
- inspiratory flow, and dynamic compliance, relationship (letter), 88:465
- protection, during endobronchial laser surgery, 88:298
- pulmonary function, after cardiac and thoracic surgery, 88:1272
- pulmonary hypertension, during minimally invasive thoracolumboendoscopic spine surgery, 88:1244

**Lung volume reduction**, see Surgery

**Lysophosphatidic acid**, see Receptors

**MAC**, see Potency, anesthetic

**Macintosh blade**, see Equipment

**Macroglossia**, see Complications

**Magnesium**, see Ions

**Magnetic resonance imaging**, see Imaging techniques

**Malignant hyperthermia susceptibility**, see Complications

**Mask induction**, see Anesthetic techniques

**Mastectomy**, see Surgery

**Maxillofacial surgery**, see Surgery

**Measurement techniques**

- Amsterdam Preoperative Anxiety and Information Scale, as measure of parents' preoperative anxiety and anesthesia knowledge, validation, 88:251
- audiometry, pure tone and speech, effects of midazolam, human volunteers, 88:1064
- brachial plexus depth, indirect vs direct measurement, 88:1113
- capnography, and laryngotracheal topical anesthesia kit, for difficult endotracheal intubation, 88:943
- cold pressor test, pain suppression by millimeter waves, 88:836
- composite sedation, remifentanyl during regional anesthesia, 88:134
- computer-assisted auscultation, effect of halothane on amplitude and frequency characteristics of heart sounds, children, 88:263

computer-assisted image analysis, erythema development after experimental thermal injury to human skin, effect of postburn topical EMLA, 88:1131

D-dimer assay, latex agglutination, detection of excess fibrinolysis, cardiac and noncardiac thoracic surgery, 88:1226

formalin test, anesthetics and spinal Fos, 88:414

gastric tonometry, thoracic epidural anesthesia and visceral perfusion and metabolism, 88:402

*in vitro* contracture test, diagnosis of malignant hyperthermia susceptibility, humans, 88:1155

laser Doppler fluxmetry, assessment of changes in rat paw perfusion after experimental mononeuropathy, 88:103

near infrared spectroscopy, continuous measurement of cerebral oxygenation during induction of anesthesia, 88:554

phonocardiography, effect of halothane on amplitude and frequency characteristics of heart sounds, children, 88:263

posttetracic count, intense neuromuscular block, anesthesia during endolaryngeal procedures, costs due to waiting time, 88:1335

quality of recovery score, after general anesthesia and surgery, development and psychometric testing, adults, 88:83

scanning electron microscopy, observations, lumbar dura mater biomechanics, 88:1317

SONOCLOTT analysis, effects of *in vitro* hemodilution with gelatin, hydroxyethyl starch, and lactated Ringer's solution on markers of coagulation, 88:483

Standard Anesthesia Learning Test, as measure of parent's preoperative anxiety and anesthesia knowledge, validation, 88:251

stimulus intensity, interactions among  $\mu$ -opioids, 88:407

transcranial Doppler ultrasonography, dynamic cerebral autoregulation, sevoflurane vs isoflurane, 88:341

visual analog scale, statistical analysis of total labor pain, and application to studies of analgesic effectiveness during childbirth (letter), 88:1193

- voltammetry, activation of rostral ventrolateral medulla during strychnine allodynia, 88:1125
- Mechanisms of anesthetic action**, see Theories of anesthetic action
- MEDLINE-based analysis**, see Statistics
- Meperidine**, see Analgesics; Anesthetics, local
- Mepivacaine**, see Anesthetics, local
- Meta-analysis**, see Statistics
- Metabolism**
  - Compound A, and changes in biochemical markers of renal injury, healthy volunteers, 88:437
  - morphine-3-glucuronide, CSF and plasma concentrations, before and after initiation of intracerebroventricular morphine for cancer pain management, 88:109
  - morphine-6-glucuronide, CSF and plasma concentrations, before and after initiation of intracerebroventricular morphine for cancer pain management, 88:109
  - visceral, intraoperative, effects of thoracic epidural anesthesia, 88:402
- Metal toxicity**, see Complications
- Methemoglobinemia**, see Complications
- Methylprednisolone**, see Pharmacology, steroids
- Metoclopramide**, see Pharmacology
- Meyer-Overton hypothesis**, see Theories of anesthetic action
- Midazolam**, see Anesthetics, intravenous; Pharmacology
- Middle latency auditory evoked potentials**, see Monitoring
- Midfemoral block**, see Anesthetic techniques, regional
- Millimeter waves**, see Anesthetic techniques
- Minimum alveolar anesthetic concentration**, see Potency, anesthetic
- Mitral valve**, see Heart
- Mivacurium**, see Neuromuscular relaxants
- Monitored anesthesia care**, see Anesthetic techniques
- Monitoring**
  - blood pressure, profound myocardial depression during acute bupivacaine overdose, pigs, 88:1143
  - capnography
    - assessment of tracheal tube position, 88:766
    - during neurosurgery, reliability in relation to various intraoperative positions, 88:43
    - endobronchial intubation, as cause of immediate increase in peak inflation pressure, pediatric patients, 88:268
  - cerebral oxygenation, after traumatic brain injury, monitoring, measuring brain tissue oxygenation vs jugular venous oxygen saturation, 88:549
  - computed tomographic scans, measurements of left main bronchus diameter, predicting size of double-lumen endobronchial tube, 88:302
  - continuous esophageal aortic blood flow echo Doppler, assessment of hemodynamic effects of pneumoperitoneum during laparoscopic surgery, healthy infants (letter), 88:468
  - electrocardiography
    - profound myocardial depression during acute bupivacaine overdose, pigs, 88:1143
    - T-wave amplitude, as guide for detecting intravascular injection of test dose, anesthetized children, 88:754
  - electrocorticography, intraoperative localization of epileptogenic focus, use of alfentanil and fentanyl, 88:1101
  - electroencephalography (EEG)
    - Bispectral Index, as measure of depth of sleep, 88:659
    - dexmedetomidine and anesthetic depth, 88:625
    - prediction of movement to noxious stimulation during 1 MAC isoflurane/N<sub>2</sub>O anesthesia, 88:1142
  - epicardial echocardiography (EE), intraoperative management of pediatric patient undergoing cardiac tumor resection, 88:104888:1048
  - esophageal, assessment of tracheal tube position, 88:766
  - Holter, after vascular surgery, high-risk patients, effect of heart rate control on myocardial ischemia, 88:477
  - intraarterial blood gas, continuous, reliability during one-lung ventilation (letter), 88:1192
  - jugular venous oxygen saturation, vs measuring brain tissue oxygenation, for monitoring cerebral oxygenation after traumatic brain injury, 88:549
  - middle latency auditory evoked potentials, and prediction of movement to noxious stimulation during 1 MAC isoflurane/N<sub>2</sub>O anesthesia, 88:1142
  - multipulse transcranial electrical stimulation, myogenic motor-evoked potentials, within-patient variability, during two levels of partial neuromuscular blockade, aortic surgery, 88:22
  - neuromuscular, common peroneal nerve stimulation, evaluation, awake volunteers and anesthetized patients, 88:187
  - neuromuscular blockade, at larynx, assessing, effect of change in resting cuff pressure and comparison with video imaging, anesthetized humans, 88:426
  - precordial stethoscope, heart sounds, amplitude and frequency characteristics, effect of halothane, children, 88:263
  - pulmonary function, changes, after interscalene brachial plexus anesthesia with 0.5% and 0.75% ropivacaine, double-blind comparison with 2% mepivacaine, 88:587
  - respiratory gas, during high-frequency jet ventilation for tracheal resection, use of double-lumen jet catheter, 88:224
  - sensory level, after subarachnoid anesthesia, assessment using pressure palpator, 88:398
  - somatosensory evoked potentials, used to compare effects of three asymmetric sternal retractors on brachial plexus function, 88:292
  - temperature
    - during neuraxial anesthesia (editorial), 88:243
    - during regional anesthesia, 88:373
  - thromboelastography, -guided transfusion algorithm, effectiveness in reducing transfusion requirements, complex cardiac surgery, 88:312
  - transcranial motor evoked potentials, electrical and magnetic transcranial stimulation during N<sub>2</sub>O/opioid anesthesia, 88:568
  - transcutaneous
    - of CO<sub>2</sub> tension, after cardi thoracic surgery, infants and children, 88:531
  - PO<sub>2</sub>, synergistic effect of sympathectomy and hyperbaric oxygen exposure, healthy volunteers, 88:67
  - transesophageal echocardiography (TEE)
    - detection of gas embolism and cardiac valvular dysfunction during laparoscopic nephrectomy, 88:500
    - intraoperative, damage to pilot cuff tubing (letter), 88:1187



**Monitoring—continued**

- intraoperative, management of
  - pediatric patient undergoing cardiac tumor resection, 88:104888:1048
- postoperative macroglossia, rapid resolution after extubation of trachea, 88:220
- systematic examination in mitral valve repair, evolution of discipline into twenty-first century (editorial), 88:1197
- systemic examination, improved evaluation of location and mechanism of mitral valve regurgitation, 88:1205
- transnasal, modified application mode for cardiac examination, ventilated patients, 88:306
- transillumination, assessment of tracheal tube position, 88:766
- Monte-Carlo computer simulation**, operating room efficiency, decrease in case duration required to complete an additional case, 88:72
- Mood**, effects of subanesthetic concentrations of sevoflurane and N<sub>2</sub>O, alone and in combination, healthy volunteers, 88:1149
- Morbidity**, anesthetic, risk of, patients with malignant osteopetrosis, 88:1292
- Morphine**, see Analgesics; Pharmacology
- Mortality**
  - anesthetic, risk of, patients with malignant osteopetrosis, 88:1292
  - postoperative, CABG, drug therapy with nitrates and  $\beta$ -adrenergic blockers as predictors of mortality and survival, 88:286
- Motoneurons**, see Brain
- Multipulse transcranial electrical stimulation**, see Monitoring
- Murphy eye**, see Equipment, endobronchial tube
- Muscle**, smooth, tracheal, tone, direct effects of heparin and protamine, dogs, 88:1232
- Myocardial depression**, see Heart
- Myringotomy**, see Surgery
- Nalbuphine**, see Analgesics
- Naloxone**, see Antagonists, opioid; Receptors, opioid, antagonists
- National Institute of Occupational Safety and Health**, recommended exposure limits for desflurane and isoflurane (letter), 88:1424
- Near infrared spectroscopy**, see Measurement techniques
- Neck**, see Position
- Needles**, see Equipment

**Neostigmine**, see Antagonists, neuromuscular relaxants

**Nephrectomy**, see Surgery

**Nephrotoxicity**, see Toxicity

**Nerve**

- brachial plexus, injury, and asymmetric retractors, 88:292
- common peroneal, stimulation, for neuromuscular monitoring, evaluation, awake volunteers and anesthetized patients, 88:197
- sciatic, new lateral approach, midfemoral block (letter), 88:1426
- sympathetic innervation of forelimb resistance vessels, cats, 88:320
- ulnar, neuromuscular monitoring, 88:197
- Nerve injury**, see Complications
- Neuraxial anesthesia**, see Anesthetic techniques
- Neurological complications**, see Complications
- Neuromuscular blockade**, see Monitoring
- Neuromuscular relaxants**
  - atracurium, intense neuromuscular block for anesthesia during endolaryngeal procedures, costs due to waiting time, 88:1335
  - cisatracurium, severe anaphylactic reaction to, 88:462
  - mivacurium
    - induced neuromuscular blockade, spontaneous recovery from, nature of, 88:648
  - intense neuromuscular block for anesthesia during endolaryngeal procedures, costs due to waiting time, 88:1335
  - resistance to, rabbits treated with chronic isepamicin, 88:654
- rapacurium (Org 9487), and its 3-desacetyl metabolite (Org 9488), pharmacokinetic-dynamic relationship, 88:640
- rocuronium
  - assessment of neuromuscular block at larynx, 88:426
  - induced neuromuscular blockade, effects of dexmedetomidine, human volunteers, 88:633
- injection, withdrawal associated with, preventing, use of lidocaine, children and adolescents, 88:746
- intense neuromuscular block for anesthesia during endolaryngeal procedures, costs due to waiting time, 88:1335
- resistance to, rabbits treated with chronic isepamicin, rabbits, 88:654

- succinylcholine, and massive intraoperative pulmonary embolism (letter), 88:1423
- suxamethonium, intense neuromuscular block for anesthesia during endolaryngeal procedures, costs due to waiting time, 88:1335
- vecuronium
  - common peroneal nerve stimulation for neuromuscular monitoring, evaluation, awake volunteers and anesthetized patients, 88:187
- intense neuromuscular block for anesthesia during endolaryngeal procedures, costs due to waiting time, 88:1335
- Neurons**, activity, nucleus raphe magnus, effect of  $\alpha$ -adrenergic drugs, rats, 88:459
- Neuropathic pain**, see Pain
- Neurosurgery**, see Surgery
- Neurosurgical anesthesia**, see Anesthesia
- Neurotoxicity**, see Toxicity
- Neurotransmitters**
  - nitric oxide, synthesis inhibition, modification of cardiotoxicity of tetracaine and lidocaine, 88:717
  - spinal cord synaptosomes, glutamate release, effects of clonidine and phorbol acetate, rats, 88:1401
- Neutrophils**, see Cells
- Nicardipine**, see Pharmacology
- Nicotinic acetylcholine receptors**, see Receptors
- Nitrates**, see Risk
- Nitric oxide**, see Gases, nonanesthetic; Neurotransmitters
- Nitroglycerin**, see Pharmacology
- Nitroprusside**, see Pharmacology
- Nitrous oxide**, see Anesthetics, gases
- N-methyl-D-aspartate**, see Receptors, NMDA
- Nociception**, see Pain
- Noncardiac surgery**, see Surgery
- Nonimmobilizers**, convulsant inhaled compounds, 88:884
- Nonsteroidal antiinflammatory drugs**
  - ketoprofen, intravenous, effect on postoperative epidural sufentanil analgesia, children, 88:1036
  - tenoxicam, effect on intraperitoneal adhesions and prostaglandin E<sub>2</sub> levels, mice, 88:939
- Norepinephrine**, see Sympathetic nervous system, catecholamines
- North American Malignant Hyperthermia Group**, protocols for diagnosis of malignant hyperthermia susceptibility, humans, 88:1155



# **Nose**

- cancerous, patient with, use of LMA before tracheal intubation (letter), 88:1429
- fracture, trigeminovagal reflex during repair under general anesthesia, 88:1183

# **Occupational exposure, see Toxicity**

# **Ondansetron, see Pharmacology**

# **One-lung ventilation, see Ventilation**

# **Operating rooms (OR)**

- efficiency
  - decrease in case duration required to complete an additional case, computer simulation study, 88:72
  - improving, at academic institutions, successful strategies (letter), 88:963
- impact of rapid anesthetic recovery protocols on ASC staffing, 88:1053

# **Opioids, see Analgesics**

# **Org 9487, see Neuromuscular relaxants, rapacuronium**

# **Orthopedic surgery, see Surgery**

# **Osmolality, see Urine**

# **Osteopetrosis, see Complications**

# **Outcome**

- cerebral ischemia, effects of aprotinin, rats, 88:1
- changing temperature management for CPB, 88:1254
- clinical anesthesia, perspective of panel of expert anesthesiologists, 88:1085
- general anesthesia and surgery, quality of recovery score, development and psychometric testing, adults, 88:83
- histopathologic, spinal cord ischemia, effects of moderate hypothermia and intrathecal tetracaine, rabbits, 88:56
- lung volume reduction surgery, preoperative functional predictors, 88:28
- neurological, spinal cord ischemia, effects of moderate hypothermia and intrathecal tetracaine, rabbits, 88:56

# **Oxycodone, see Analgesics**

# **Oxygen**

- consumption, myocardial, pacing-induced changes, effect of nitroglycerin, patients with coronary artery disease, 88:271
- delivery
  - during hyperbaric oxygen treatment, enhancement by sympathetic nerve block, 88:67
  - during retrograde cerebral perfusion, humans, 88:8
- extraction, cerebral, during CPB, effects of propofol (letter), 88:231

- hyperbaric oxygen exposure, and sympathectomy, synergistic effect on transcutaneous PO<sub>2</sub>, healthy volunteers, 88:67

# **Oxygenation**

- arterial, during one-lung ventilation, combined vs general anesthesia, 88:494
- brain tissue, vs jugular venous oxygen saturation, measuring, for monitoring cerebral oxygenation after traumatic brain injury, 88:549
- cerebral, continuous measurement by near infrared spectroscopy during induction of anesthesia, 88:554

# **Pacemakers, see Equipment**

# **Pain**

- antinociception
  - interactions among  $\mu$ -opioids, 88:407
  - morphine and monoamine antinociception, rats, 88:141
  - xenon and N<sub>2</sub>O and spinal dorsal horn neurons, 88:893
- cancer
  - anesthesia, endothelin and major surgery, 88:898
  - management, intracerebroventricular morphine, CSF and plasma concentrations of morphine, morphine-3-glucuronide, and morphine-6-glucuronide, 88:109
- erythromelalgia, primary, in upper extremities, endoscopic thoracic sympathectomy, 88:865
- hyperalgesia, secondary, reduction by adenosine, two human models of cutaneous inflammatory pain, 88:605
- inflammatory, cutaneous, secondary hyperalgesia, effect of adenosine, 88:605
- injection, rocuronium, alleviation by pretreatment with intravenous lidocaine, 88:746
- intraoperative, during postpartum tubal ligation, effect of intraperitoneal lidocaine, 88:1077
- labor
  - epidural analgesia, initiation, epidural ropivacaine, dose-finding study, 88:1340
  - statistical analysis using visual analog scale and application to studies of analgesic effectiveness during childbirth (letter), 88:1193
  - visual analog scale scores (letter), 88:1421

- low back, chronic, patients with, percutaneous electrical nerve stimulation, analgesic response, effect of stimulus frequency, 88:841

# **neuropathic**

- allodynia, strychnine-induced, activation of rostral ventrolateral medulla, rats, 88:1125
- rat paw perfusion after mononeuropathy, 88:103

# **nociception**

- adenosine reduces secondary hyperalgesia, 88:605
- effects of clonidine and phorbol acetate on glutamate release from spinal synaptosomes, rats, 88:1401
- peripheral antihyperalgesic effect of morphine to heat, but not mechanical stimulation, healthy volunteers, after ultraviolet-B irradiation, 88:117

# **postoperative**

- abdominal surgery, epidural analgesia with local anesthetics, earlier motor recovery with 0.2% ropivacaine than 0.175% bupivacaine, 88:128
- abdominal surgery, major, epidural ropivacaine infusion alone vs in combination with 1, 2, and 4  $\mu$ g/mL fentanyl, 88:857
- acute, safety and efficacy of intrathecal opioids, 88:599
- adenotonsillectomy, dextromethorphan premedication, children, 88:749
- anterior cruciate ligament surgery, postoperative analgesia with controlled-release oxycodone, 88:1286
- cesarean delivery, determination of effective dose of intrathecal morphine for pain relief, 88:346
- clonidine administered as axillary block, 88:1109
- craniotomy procedures, 88:335
- epidural sufentanil anesthesia, effect of intravenous ketoprofen, children, 88:1036
- laparoscopic cholecystectomy, effect of preincisional dextromethorphan treatment, 88:1331
- major spinal surgery, persistently high plasma morphine-6-glucuronide levels despite decreased hourly PCA morphine use after single-dose diclofenac, 88:1137
- recovery characteristics of sevoflurane and halothane, preschool-aged children undergoing bilateral

**Pain—continued**

- myringotomy and pressure equalization tube insertion, 88:34
- sameridine and spinal anesthetic, 88:815
- total hip surgery, amount of morphine needed for postoperative analgesia, and intensity of preoperative pain, direct correlation, 88:146
- total hip surgery, intrathecal morphine, dose-finding study, 88:822
- preoperative, intensity, and amount of morphine needed for postoperative analgesia, direct correlation, 88:146
- response, effects of subanesthetic concentrations of sevoflurane and  $N_2O$ , alone and in combination, healthy volunteers, 88:1149
- sensation, suppression, caused by millimeter waves, 88:836
- therapies, implantable devices
  - nationwide survey of academic teaching programs, 88:1311
  - training requirements (editorial), 88:1203
- tourniquet, prevention, adding clonidine to lidocaine for intravenous regional anesthesia, 88:1327
- Panic disorder**, see Complications
- Paranoia**, see Psychology
- Paravertebral blockade**, see Anesthetic techniques
- Paresthesia**, see Complications
- Parkinson surgery**, see Surgery
- PASS**, see Sedation, pediatric analog sedation score
- Patient-controlled analgesia**, see Anesthetic techniques
- Patient preferences**, for early discharge after laparoscopic cholecystectomy, 88:1280
- PCA**, see Anesthetic techniques, patient-controlled analgesia
- Pediatric anesthesia**, see Anesthesia
- Pentobarbital**, see Anesthetics, intravenous
- Peptides**, vasodilating, adrenomedullin, increased plasma concentration in hemorrhagic shock, dogs, 88:326
- Percutaneous electrical nerve stimulation**, see Anesthetic techniques
- Perfusion**
  - retrograde cerebral perfusion, oxygen delivery during, humans, 88:8
  - visceral, intraoperative, effects of thoracic epidural anesthesia, 88:402
- Peripheral nerve block**, see Anesthetic techniques, regional
- Perphenazine**, see Antiemetics

**Pharmacodynamics, propofol**

- concentration and LMA vs cuffed oropharyngeal airway, 88:917

**Pharmacokinetics**

- enflurane, "second gas effect" is not valid concept, 88:188
- halothane, rate of rise, children with shunt, 88:759
- morphine, persistently high plasma morphine-6-glucuronide levels despite decreased hourly PCA morphine use after single-dose diclofenac, 88:1137
- $N_2O$ , "second gas effect" is not valid concept, 88:188
- rapacurium (Org 9487) and its 3-desacetyl metabolite (Org 9488), 88:640
- vasopressin, intrinsic, effects on urinary aquaporin-2 excretion and urine osmolality during surgery under general anesthesia, 88:181

**Pharmacology**

- adenosine, reduction of secondary hyperalgesia, two human models of cutaneous inflammatory pain, 88:605
- angiotensin-converting enzyme inhibitors, patients chronically treated with, intraoperative refractory hypotension, treatment with terlipressin, 88:980
- aprotinin, effects on outcome from cerebral ischemia, rats, 88:1
- $\beta$ -blockers
  - drug therapy with, as predictor of survival after CABG, 88:286
  - heart rate control and ischemia, 88:286
  - interscalene block,  $\beta$ -blockers, and vasovagal reactions, 88:378
  - therapy, individualizing, patient-specific target-based heart rate control (editorial), 88:475
- clonidine
  - administered as axillary block, and postoperative pain, 88:1109
  - axillary block, tramadol plus mepivacaine, 88:853
  - effect on glutamate release from spinal synaptosomes, rats, 88:1401
  - effects on human neutrophil functions, 88:452
  - effects on sensitivity to phenylephrine and nitroprusside, patients with essential hypertension recovering from surgery, 88:1239
  - local anesthetic and, intravascular injection, followed by prolonged unconsciousness, reversal by naloxone, 88:1185

- spinal, prolongs labor analgesia from spinal sufentanil and bupivacaine, 88:573
- vs high thoracic epidural anesthesia, perioperative stress response, patients undergoing CABG, 88:701
- desipramine
  - antinociceptive effects after intrathecal injection, rats, 88:141
  - combined with morphine, antinociceptive effects after intrathecal injection, rats, 88:141
  - combined with serotonin, antinociceptive effects after intrathecal injection, rats, 88:141
- dexamethasone, granisetron/dexamethasone combination, for reducing nausea and vomiting during and after spinal anesthesia for cesarean section, 88:1346
- dexmedetomidine, effects on human neutrophil functions, 88:452
- dextromethorphan, preincisional, effect on postoperative pain and opioid requirement after laparoscopic cholecystectomy, 88:1331
- diclofenac, does not decrease renal blood flow or glomerular filtration, elderly patients undergoing orthopedic surgery, 88:149
- droperidol
  - efficacy and adverse effects during PCA, quantitative systematic review, 88:1354
  - efficacy and safety for preventing PONV, meta-analysis, 88:1370
  - vs ondansetron, for PONV (letter), 88:465
- endothelin-1 (ET-1), perioperative plasma endothelin-1 and Big ET-1, elderly patients undergoing major surgical procedures, 88:898
- enoxaparin, and epidural hematomas (letter), 88:229
- epinephrine, and diagnostic accuracy of aspiration during labor epidural analgesia, 88:1073
- granisetron
  - granisetron/dexamethasone combination, for reducing nausea and vomiting during and after spinal anesthesia for cesarean section, 88:1346
  - vs perphenazine, for reducing PONV after tonsillectomy, children, 88:1298
- $H_2$  blocking drugs, for acid aspiration prophylaxis, and anesthesia for cesarean section, German survey, 88:63

insulin, continuous infusion, and neutrophil function, diabetic cardiac surgery patients, 88:1011

isoproterenol  
-containing test dose, effective dose, anesthetized children (letter), 88:1189  
epidural analgesia and (letter), 88:1190  
labeling, and label, terminology not interchangeable (letter), 88:467

metoclopramide  
efficacy and safety for preventing PONV, meta-analysis, 88:1370  
repetitive peridural administration, histopathological changes after, dogs, 88:100

midazolam, neurotoxicity of spinal drugs, 88:797

morphine  
antinociceptive effects after intrathecal injection, rats, 88:141  
combined with desipramine, antinociceptive effects after intrathecal injection, rats, 88:141  
combined with serotonin, antinociceptive effects after intrathecal injection, rats, 88:141

nicardipine, attenuation of hemodynamic aberrances during laparoscopic adrenalectomy for pheochromocytoma, 88:16

nitroglycerin, effect on pacing-induced changes in myocardial oxygen consumption and metabolic coronary vasodilation, patients with coronary artery disease, 88:271

nitroprusside  
reduction of depressant effects of protamine sulfate on isolated ischemic rat hearts, 88:710  
sensitivity to, patients with essential hypertension recovering from surgery, effects of clonidine, 88:1239

ondansetron  
efficacy and safety for preventing PONV, meta-analysis, 88:1370  
vs droperidol, for PONV (letter), 88:465

perphenazine, vs granisetron, for reducing PONV after tonsillectomy, children, 88:1298

phenylephrine  
effect on hypotension induced by alkalized lidocaine epidural anesthesia, 88:1322  
sensitivity to, patients with essential hypertension recovering from surgery, effects of clonidine, 88:1239

phorbol acetate, effect on glutamate release from spinal synaptosomes, rats, 88:1401

prostaglandin E<sub>1</sub>, effects on intraoperative temperature changes and incidence of postoperative shivering during deliberate mild hypothermia for neurosurgical procedures, 88:446

protamine  
administration, catastrophic pulmonary vasoconstriction arising from, managing, use of NO, 88:505  
depressant effects on isolated ischemic rat hearts, reduced by isoflurane and sodium nitroprusside, 88:710

serotonin  
antinociceptive effects after intrathecal injection, rats, 88:141  
combined with desipramine, antinociceptive effects after intrathecal injection, rats, 88:141  
combined with morphine, antinociceptive effects after intrathecal injection, rats, 88:141  
sodium citrate, for acid aspiration prophylaxis, and anesthesia for cesarean section, German survey, 88:63

steroids  
epidural, treatment of failed back surgery syndrome, cost-benefit of fluoroscopy, 88:467  
epidural, treatment of failed back surgery syndrome (editorial), 88:240  
methylprednisolone, prevention of increased circulating tumor necrosis factor- $\alpha$  levels and lung injury associated with systemic inflammatory response syndrome due to intraperitoneal hyperthermia, 88:771

strychnine, -induced allodynia, activation of rostral ventrolateral medulla, rats, 88:1125

terlipressin, treatment of intraoperative refractory hypotension, patients chronically treated with antagonist of renin-angiotensin system, 88:980

vasopressin  
effects on endotoxin-induced attenuation of contractile responses, human gastroepiploic arteries *in vitro*, 88:542  
post-CPB refractory hypotension and (letter), 88:695

**Phenylephrine**, see Pharmacology  
**Pheochromocytoma**, see Complications  
**Phonocardiography**, see Measurement techniques

**Phorbol acetate**, see Pharmacology  
**Plasma**, see Blood  
**Plastic surgery**, see Surgery  
**Platelets**, see Coagulation  
**Pneumocephalus**, see Complications  
**Pneumoperitoneum**, see Carbon dioxide, insufflation

**Polyhalogenated alkanes**  
convulsant inhaled compounds, 88:884  
1,2-dichlorohexafluorocyclobutane, effect on depolarization-evoked glutamate release, rat cortical brain slices, 88:1168  
1,1,2-trichlorotrifluoroethane, effect on depolarization-evoked glutamate release, rat cortical brain slices, 88:1168

**Polyhalogenated methyl ethyl ethers**, see Anesthetics, volatile

**Position**  
anatomic, disposable vs reusable LMA, 88:1418  
head, influence of oropharyngeal leak pressure and cuff position with flexible and standard LMA, 88:913  
intraoperative, and reliability of capnography monitoring during neurosurgery, 88:43  
neck, influence of oropharyngeal leak pressure and cuff position with flexible and standard LMA, 88:913  
sitting, shoulder surgery, under interscalene block, vasovagal episodes and  $\beta$ -adrenergic blockers, 88:378

**Positive end-expiratory pressure**, see Ventilation

**Postanesthesia care units (PACU)**, impact of rapid anesthetic recovery protocols on ASC staffing, 88:1053

**Posttétanic count**, see Measurement techniques

**Potency, anesthetic**, minimum alveolar anesthetic concentration (MAC)  
dexmedetomidine and anesthetic depth, 88:625  
fluorinated alkanols, relevance to theories of narcosis, rats, 88:867  
intravenous halothane anesthesia, 88:671  
1 MAC isoflurane/N<sub>2</sub>O anesthesia, movement to noxious stimulation during, prediction, middle latency auditory evoked responses and EEG-derived variables, 88:114288:1412  
polyhalogenated methyl ethyl ethers, 88:1161

- Potency, anesthetic**, minimum alveolar anesthetic concentration (MAC)—*continued*  
 2 MAC isoflurane, cesarean delivery of twin gestation, one normal and one with large neck mass, 88:594  
 volatile anesthetics  
 normal and cardiomyopathic hamsters, 88:489  
 production of immobility by acting on two sites approximately five carbon atoms apart, 88:1395
- Precordial stethoscope**, see **Monitoring**
- Preeclampsia**, see **Complications**
- Preemptive analgesia**, see **Anesthetic techniques**
- Premedication**, dextromethorphan, oral, does not reduce pain or analgesic consumption after adenotonsillectomy, children, 88:749
- Preoperative**, anxiety  
 distress during induction of anesthesia, and postoperative behavioral outcomes, 88:1042  
 parental  
 and anesthesia knowledge, measures, validation, 88:251  
 perioperative information (editorial), 88:237  
 use of preanesthetic video before pediatric ambulatory surgery, 88:246
- Pressor sites**, see **Central nervous system**
- Pressure palpator**, see **Equipment**
- Prolonged unconsciousness**, see **Complications**
- Propacetamol**, see **Analgesics**
- Propofol**, see **Anesthetics**, intravenous
- Prostaglandin E<sub>1</sub>**, see **Pharmacology**
- Prostaglandin E<sub>2</sub>** levels, effect of tenoxicam, mice, 88:939
- Protamine**, see **Coagulation; Pharmacology**
- Protein S deficiency**, see **Complications**
- Psychological testing**, predictive evaluation of residents, 88:1092
- Psychology**, paranoia, emergence "delirium" after sevoflurane anesthesia, 88:1308
- Psychomotor performance**, effects of subanesthetic concentrations of sevoflurane and N<sub>2</sub>O, alone and in combination, healthy volunteers, 88:1149
- Publications**, important anesthesia and critical care journals, national origin of articles, MEDLINE-based analysis, 88:1175
- Pulmonary embolism**, see **Complications**
- Pulmonary function**, see **Monitoring**
- Pulmonary hypertension**, see **Complications; Lung**
- Pulse oximeter**, see **Equipment**
- Pupillary dilatation**, see **Eyes**
- Quality of recovery score**, see **Measurement techniques**
- Rapacuronium**, see **Neuromuscular relaxants**
- Raphe magnus**, see **Brain**
- Receptors**  
 $\alpha$ -adrenergic  
 agonists, and neutrophil functions, 88:452  
 agonists, dexmedetomidine, effects on neuromuscular blockade, human volunteers, 88:633  
 agonists, effect on activity of neurons, nucleus raphe magnus, rats, 88:459  
 antagonists, effect on activity of neurons, nucleus raphe magnus, rats, 88:459  
 $\alpha_2$ -adrenergic, agonists, clonidine, added to lidocaine, for intravenous regional anesthesia, prevention of tourniquet pain, 88:1327  
 $\beta$ -adrenergic, antagonists  
 heart rate control and ischemia, 88:477  
 patient-specific target-based heart rate control (editorial), 88:475  
 GABA, general anesthetics and synaptic transmission, 88:676  
 GABA<sub>A</sub>, actions of fluorinated alkanols, relevance to theories of narcosis, 88:877  
 lysophosphatidic acid, signaling, synergistic inhibition by charged and uncharged local anesthetics, 88:1117  
 nicotinic acetylcholine  
 neuronal, inhibition by depressant and convulsant barbiturates, 88:1406  
 P<sub>2X</sub> purinoceptors and, in PC12 cells, effects of ketamine and propofol, 88:174  
 NMDA  
 antagonists, dextromethorphan, premedication, 88:754  
 general anesthetics and synaptic transmission, 88:676  
 non-NMDA, general anesthetics and synaptic transmission, 88:676  
 opioid, antagonists, naloxone, inhibition of gastric emptying, rats, 88:204
- Recovery**  
 awakening, comparison of induction methods, 88:930  
 fast-tracking, after outpatient anesthesia, new criteria, comparison with modified Aldrete's scoring system, 88:1069
- from sevoflurane and halothane, preschool-aged children undergoing bilateral myringotomy and pressure equalization tube insertion, 88:34  
 impact of rapid anesthetic recovery protocols on ASC staffing, 88:1053  
 quality of recovery score, after general anesthesia and surgery, development and psychometric testing, adults, 88:83  
 spontaneous, from mivacurium-induced neuromuscular block, nature of, 88:648
- Reflexes**, trigeminovagal reflex, during repair of nasal fracture under general anesthesia, 88:1183
- Reflex sympathetic dystrophy**, see **Complications**
- Regional anesthesia**, see **Anesthetic techniques**
- Regional anesthesia and pain management**, see **Anesthesia**
- Regurgitation**, see **Complications**
- Remifentanyl**, see **Analgesics; Anesthetics**, intravenous
- Renal injury**, see **Kidney**
- Residents**, see **Education**
- Respiratory arrest**, see **Complications**
- Reticular formation**, see **Brain**
- Retinal artery air embolism**, see **Eye**
- Right-to-left shunt**, see **Heart**
- Ringer's lactate**, see **Fluid balance**, lactated Ringer's solution
- Risk**  
 $\beta$ -blockers, drug therapy with, as predictor of survival after CABG, 88:286  
 factors, tracheal intubation without muscle relaxants, 88:737  
 metal toxicity, in combined spinal-epidural anesthesia, 88:393  
 morbidity and mortality, anesthetic, patients with malignant osteopetrosis, 88:1292  
 nitrates, drug therapy with, as predictor of mortality after CABG, 88:286  
 persistent paresthesia, and repeated axillary block, 88:382
- Rocuronium**, see **Neuromuscular relaxants**
- Ropivacaine**, see **Anesthetics**, local
- Rostral ventrolateral medulla**, see **Brain**
- Samerdine**, see **Analgesics; Anesthetics**, local
- Scanning electron microscopy**, see **Measurement techniques**
- Sciatic nerve**, see **Nerve**
- Scoring systems**  
 modified Aldrete's scoring system, comparison with new criteria for



- fast-tracking after outpatient anesthesia, 88:1069
- pediatric analog sedation score (PASS), simple (letter), 88:227
- quality of recovery score, after general anesthesia and surgery, development and psychometric testing, adults, 88:83
- visual analog scale scores for labor pain (letter), 88:1421
- Sedation**
  - during monitored anesthesia care, cost-effectiveness of methohexital vs propofol, 88:723
  - pediatric analog sedation score (PASS), simple (letter), 88:227
  - remifentanyl during regional anesthesia, 88:134
  - resting ventilation, effects of intravenous infusions of morphine or sameridine, 88:160
- Sensory level**, see Monitoring
- Serotonin**, see Pharmacology
- Sevoflurane**, see Anesthetics, volatile
- Shivering**, see Complications
- Shock**, see Complications
- Shoulder surgery**, see Surgery
- Sitting position**, see Position
- Sleep**, depth of, Bispectral Index as measure, 88:659
- Smooth muscle**, see Muscle
- Sodium chloride**, see Fluid balance
- Sodium citrate**, see Pharmacology
- Somatosensory evoked potentials**, see Monitoring
- Spinal analgesia**, see Anesthetic techniques
- Spinal cord**
  - blood flow, age-related effects of
    - epidural lidocaine, with and without epinephrine, anesthetized rabbits, 88:1302
  - descending inhibitory system, xenon and N<sub>2</sub>O and spinal dorsal horn neurons, 88:893
  - formalin-induced Fos immunoreactivity, pre- vs postinjury effects of intravenous GABAergic anesthetics, rats, 88:414
  - ischemia, transient, effects of moderate hypothermia and intrathecal tetracaine, rabbits, 88:56
  - synaptosomes, glutamate release, effects of clonidine and phorbol acetate, rats, 88:1401
- Spinal cord stimulation**, see Anesthetic techniques
- Spinal surgery**, see Surgery, back
- Standard Anesthesia Learning Test**, see Measurement techniques
- Statistics**
  - analysis of total labor pain using visual analog scale and application to studies of analgesic effectiveness during childbirth (letter), 88:1193
  - MEDLINE-based analysis, important anesthesia and critical care journals, national origin of articles, 88:1175
  - meta-analysis
    - comparative efficacy and safety of ondansetron, droperidol, and metoclopramide for preventing PONV, 88:1370
    - efficacy and adverse effects of prophylactic antiemetics during PCA, 88:1354
    - use, effect on understanding of therapies for PONV (editorial), 88:1200
    - use of nonpharmacological techniques to prevent PONV, 88:1362
- Stellate ganglion**, sympathetic innervation of forelimb resistance vessels, cats, 88:320
- Steroids**, see Pharmacology
- Stimulus intensity**, see Measurement techniques
- Strychnine**, see Pharmacology
- Succinylcholine**, see Neuromuscular relaxants
- Sufentanil**, see Analgesics
- Supraclavicular nerve block**, see Anesthetic techniques
- Surgery**
  - abdominal
    - epidural analgesia with local anesthetics after, earlier motor recovery with 0.2% ropivacaine than 0.175% bupivacaine, 88:128
    - major, postoperative analgesia, epidural ropivacaine infusion alone vs in combination with 1, 2, and 4 µg/mL fentanyl, 88:857
    - upper, early postoperative hypoxemia, 88:213
  - adenotonsillectomy, postoperative analgesia, and preoperative oral dextromethorphan, children, 88:749
  - adrenalectomy, laparoscopic, for pheochromocytoma, hemodynamic changes and catecholamine release during, 88:16
  - alcohol withdrawal syndrome, prevention and treatment, 88:946
  - ambulatory, discharge criteria and complications after, 88:508
  - aortic
    - reconstruction, oxygen delivery during retrograde cerebral perfusion, humans, 88:
- two levels of partial neuromuscular blockade, within-patient variability of myogenic motor-evoked potentials to multipulse transcranial electrical stimulation, 88:22
- back**
  - failed, treatment, use of epidural steroids, cost-benefit of fluoroscopy, 88:467
  - failed, treatment, use of epidural steroids (editorial), 88:240
  - major spinal surgery, persistently high plasma morphine-6-glucuronide levels despite decreased hourly PCA morphine use after single-dose diclofenac, 88:1137
  - thoraco-lumboendoscopic spine surgery, minimally invasive, cardiopulmonary dysfunction during, 88:1244
- cardiac**
  - asymmetric retractors and nerve injury, 88:292
  - complex, transfusion requirements, reduction, effectiveness of thromboelastography-guided transfusion algorithm, 88:312
  - D-dimer formation during, 88:1226
  - heparin and protamine effects on tracheal smooth muscle, 88:1232
  - insulin infusion, and neutrophil function, diabetic patients, 88:1011
  - mitral valve regurgitation, evaluation of location and mechanism, improved, with systematic TEE examination, 88:1205
  - mitral valve repair, systematic TEE examination, evolution into twenty-first century (editorial), 88:1197
  - nitroglycerin and metabolic coronary flow control, 88:271
  - platelet function, postoperative, effect of intravenously administered magnesium, 88:1213
  - pulmonary function, postoperative, 88:1272
  - tumor resection, intraoperative management with aid of TEE and EE, pediatric patient, 88:1048
- cardiopulmonary bypass (CPB)**
  - cerebral blood flow velocity and cerebral oxygen extraction, effects of propofol (letter), 88:231
  - changing temperature management, 88:1254
  - hypotension, postoperative, and vasopressin (letter), 88:695



**Surgery—continued**

- hypothermic, cerebral production of adrenomedullin after, adult cardiac surgical patients, 88:1030
- intracranial pressure and venous cannulation, 88:329
- oxygen delivery during retrograde cerebral perfusion, humans, 88:8
- platelet hyporeactivity, young infants, 88:258
- tissue heat content and distribution, 88:1220
- cardiothoracic, transcutaneous monitoring of CO<sub>2</sub> tension after, infants and children, 88:531
- cesarean section
  - anesthesia for, and acid aspiration prophylaxis, German survey, 88:63
  - epidural analgesia and, relationship (letter), 88:963
  - epidural anesthesia, accidental dural puncture, pneumocephalus with intense headache and unilateral pupillary dilatation after, 88:582
  - epidural anesthesia, patient with von Hippel-Lindau disease and multiple sclerosis, 88:1083
  - nausea and vomiting during and after spinal anesthesia, reducing, efficacy of granisetron/dexamethasone combination, 88:1346
  - postoperative analgesia, determination of effective dose of intrathecal morphine, 88:346
  - twin gestation under 2 MAC isoflurane, one normal and one with large neck mass, 88:584
- cholecystectomy, laparoscopic
  - early discharge after, patient preferences, 88:1280
  - postoperative pain and opioid requirement, effect of preincisional dextromethorphan treatment, 88:1331
- coronary artery bypass grafting (CABG)
  - mortality and survival, postoperative, drug therapy with nitrates and  $\beta$ -adrenergic blockers as predictors, 88:286
  - perioperative stress response and release of troponin T, high thoracic epidural anesthesia vs clonidine, 88:701
- craniotomy procedures, analgesic requirements, 88:335
- endolaryngeal procedures, intense neuromuscular block, costs due to waiting time, 88:1335
- epilepsy, intraoperative localization of epileptogenic focus, use of alfentanil and fentanyl, 88:1101
- gynecologic, postoperative analgesia, propacetamol vs ketorolac in combination with PCA morphine, efficacy and tolerability, 88:611
- hernia repair, spinal anesthesia, sameridine, safety and efficacy, 88:815
- hip, total
  - intensity of preoperative pain, and amount of morphine needed for postoperative analgesia, direct correlation, 88:146
  - postoperative analgesia, intrathecal morphine, dose-finding study, 88:822
- intranasal, tracheal intubation vs flexible reinforced LMA, 88:421
- knee
  - anterior cruciate ligament, outpatient, postoperative analgesia with controlled-release oxycodone, 88:1286
  - arthroscopic meniscectomy, outpatient, postoperative analgesia with intraarticular clonidine, 88:729
- laparoscopic
  - adrenalectomy, for pheochromocytoma, hemodynamic changes and catecholamine release during, 88:16
  - cholecystectomy, early discharge after, patient preferences, 88:1280
  - cholecystectomy, postoperative pain and opioid requirement, effect of preincisional dextromethorphan treatment, 88:1331
  - nephrectomy, gas embolism and cardiac valvular dysfunction during, detection by TEE, 88:500
  - pneumoperitoneum, hemodynamic effects, assessment by continuous esophageal aortic blood flow echo Doppler, healthy infants (letter), 88:468
- laser, endobronchial, anesthesia, modified technique, 88:298
- liver transplantation, orthotopic, transfusion requirements (letter), 88:692
- lung volume reduction
  - acute myocardial infarction during, 88:332
  - preoperative functional predictors for postoperative outcome, 88:28
- mastectomy, modified radical, paravertebral blockade, pregnant patient, 88:1351
- maxillofacial, nasal fracture repair, under general anesthesia, trigeminovagal reflex during, 88:1183
- myringotomy, bilateral, and pressure equalization tube insertion, recovery characteristics of sevoflurane and halothane, preschool-aged children, 88:34
- nephrectomy, laparoscopic, gas embolism and cardiac valvular dysfunction during, detection by TEE, 88:500
- neurosurgery, capnography monitoring during, reliability in relation to various intraoperative positions, 88:43
- noncardiac, thoracic surgery, D-dimer formation during, 88:1226
- orthopedic, elderly patients undergoing, diclofenac does not decrease renal blood flow or glomerular filtration, 88:149
- Parkinson surgery, venous air embolism during, patients with spontaneous ventilation, 88:793
- plastic, elective superficial, early postoperative hypoxemia, 88:213
- shoulder
  - quadriplegic patient undergoing, autonomic hyperreflexia, prevention, use of interscalene regional anesthesia, 88:98
  - sitting position under interscalene block,  $\beta$ -adrenergic blockers and vasovagal episodes, 88:378
- thoracic
  - cardiac and noncardiac, D-dimer formation during, 88:1226
  - pulmonary function, postoperative, 88:1272
  - thoracoabdominal, early postoperative hypoxemia, 88:213
  - thoracoscopic, staging, bronchial rupture by double-lumen endobronchial tube, 88:1252
  - thoracostomy tube removal, pain relief during, EMLA vs intravenous morphine, 88:1107
  - tonsillectomy, PONV, reducing, use of preoperative oral antiemetics, granisetron vs perphenazine, children, 88:1298
  - tracheal resection, high-frequency jet ventilation, respiratory gas

- monitoring using double-lumen jet catheter, 88:224
- tubal ligation, postpartum, intraoperative pain, effect of intraperitoneal lidocaine, 88:1077
- vascular, myocardial ischemia, postoperative, effect of heart rate control, high-risk patients, 88:477
- Suxamethonium**, see Neuromuscular relaxants
- Sympathectomy**, see Anesthetic techniques
- Sympathetic nervous system** catecholamines
  - epinephrine, brain and plasma concentrations during near-complete cerebral ischemia, effects of isoflurane, ketamine, and fentanyl/N<sub>2</sub>O, rats, 88:787
  - epinephrine, plasma concentrations, changes, during laparoscopic adrenalectomy for pheochromocytoma, 88:16
  - epinephrine, with epidural lidocaine, age-related effects on spinal cord blood flow, anesthetized rabbits, 88:1302
  - norepinephrine, brain and plasma concentrations during near-complete cerebral ischemia, effects of isoflurane, ketamine, and fentanyl/N<sub>2</sub>O, rats, 88:787
  - norepinephrine, plasma concentrations, changes, during laparoscopic adrenalectomy for pheochromocytoma, 88:16
  - stress response to insertion of Combitube, LMA, or tracheal intubation, 88:1389
  - sympathetic innervation of forelimb resistance vessels, cats, 88:320
- Synaptosomes**, see Spinal cord
- Systemic inflammatory response syndrome**, see Complications
- Tachycardia**, see Complications; Heart, rate
- TCI**, see Anesthetic techniques, target-controlled infusion
- Temperature**
  - changes, intraoperative, effects of prostaglandin E<sub>1</sub>, neurosurgical patients undergoing deliberate mild hypothermia, 88:446
  - core, sevoflurane vs propofol, 88:921
  - management, changing, for CPB, 88:1254
  - monitoring
    - and management during neuraxial anesthesia (editorial), 88:243
    - practices, during regional anesthesia, 88:373
    - tissue heat content and distribution during and after CPB, 88:1220
- Tenoxicam**, see Non-steroidal anti-inflammatory drugs
- Terlipressin**, see Pharmacology
- Tetracaine**, see Anesthetics, local
- Theories of anesthetic action**
  - barbiturates, depressant and convulsant, inhibition of neuronal nicotinic acetylcholine receptors, 88:1406
  - convulsant inhaled compounds, 88:884
  - local anesthetics, charged and uncharged, synergistic inhibition of lysophosphatidic acid signaling, 88:1117
  - mechanisms
    - actions of fluorinated alkanols on GABA<sub>A</sub> receptors, relevance to theories of narcosis, 88:877
    - $\alpha$ -adrenergic effect on raphe magnus, 88:459
    - cellular excitability, volatile anesthetics and T-type currents, 88:168
    - different effects of volatile anesthetics and polyhalogenated alkanes on depolarization-evoked glutamate release, rat cortical brain slices, 88:1168
    - MAC of fluorinated alkanols, relevance to theories of narcosis, rats, 88:867
    - spinal-epidural anesthesia, 88:810
    - volatile anesthetics, production of immobility by acting on two sites approximately five carbon atoms apart, 88:1395
- Meyer-Overton hypothesis**
  - convulsant inhaled compounds, 88:884
  - MAC of fluorinated alkanols, 88:867
  - solubilities and anesthetic properties of polyhalogenated methyl ethyl ethers, 88:1161
- Theories of convulsant action**, convulsant inhaled compounds, 88:884
- Thiopental**, see Anesthetics, intravenous
- Thoracic epidural**, see Anesthetic techniques
- Thoracic surgery**, see Surgery
- Thoracoabdominal surgery**, see Surgery
- Thoraco-lumboendoscopic spine surgery**, see Surgery
- Thoracoscopic surgery**, see Surgery
- Thoracostomy tube removal**, see Surgery
- Thromboelastography**, see Monitoring
- Tolerance**, morphine, CSF and plasma concentrations, before and after initiation of intracerebroventricular morphine for cancer pain management, 88:109
- Tonsillectomy**, see Surgery
- Topical anesthesia**, see Anesthetic techniques
- Total intravenous anesthesia**, see Anesthetic techniques
- Tourniquet pain**, see Pain
- Toxicity**
  - cardiac, bupivacaine, maintenance of blood pressure during profound myocardial depression, pigs, 88:1143
  - intravenous halothane, 88:671
  - kidney, Compound A and renal injury, humans, 88:437
  - local anesthetics, modification by nitric oxide synthesis inhibition, 88:717
  - metal, risk, in combined spinal-epidural anesthesia, 88:393
  - nephrotoxicity, sevoflurane, disclosing of data used as basis for publication of studies (letter), 88:693
  - neurotoxicity, of drugs given intrathecally, 88:797
  - occupational exposure
    - recommended exposure limits for desflurane and isoflurane (letter), 88:1424
    - to sevoflurane and N<sub>2</sub>O, 88:925
  - opioid-related, potential, persistently high plasma morphine-6-glucuronide levels despite decreased hourly PCA morphine use after single-dose diclofenac, 88:1137
  - systemic, intravenous bupivacaine, comparison of effects of propofol and sevoflurane, rats, 88:155
- Toxins**, endotoxins, -induced attenuation of contractile responses, human gastroepiploic arteries *in vitro*, 88:542
- Trachea**, smooth muscle tone, direct effects of heparin and protamine, dogs, 88:1232
- Tracheal resection**, see Surgery
- Tracheo-innominate artery fistula**, see Complications
- Tracheostomy**, see Complications
- Training courses**, see Education
- Tramadol**, see Analgesics
- Transcranial Doppler ultrasonography**, see Measurement techniques
- Transcranial motor evoked potentials**, see Monitoring
- Transcutaneous monitoring**, see Monitoring

- Transesophageal echocardiography**, see Monitoring
- Transfusions**, see Blood
- Transillumination**, see Monitoring
- Transitional compounds**, convulsant inhaled compounds, 88:884
- Transnasal TEE**, see Monitoring, transesophageal echocardiography
- Trauma**, see Complications, closed head trauma
- 1,1,2-Trichlorotrifluoroethane**, see Polyhalogenated alkanes
- Trigeminal reflex**, see Reflexes
- Tubal ligation**, see Surgery
- Tumor**, see Brain; Heart
- Tumor necrosis factor- $\alpha$** , levels, increased, associated with systemic inflammatory response syndrome due to intraperitoneal hyperthermia, prevention by methylprednisolone, 88:771
- T-wave amplitude**, see Monitoring, electrocardiography
- Ulnar nerve**, see Nerve
- Upper respiratory tract infections**, see Complications
- Urine**  
aquaporin-2 excretion, effects of intrinsic vasopressin during surgery under general anesthesia, 88:181  
osmolality, effects of intrinsic vasopressin during surgery under general anesthesia, 88:181  
urinary markers, assessment of renal effects of sevoflurane, elderly patients (letter), 88:1426
- Valvular dysfunction**, see Heart
- Vaporizers**, see Equipment
- Vascular reactivity**, clonidine effects on sensitivity to phenylephrine and nitroprusside, 88:1239
- Vasoconstriction**, effects of dexmedetomidine on neuromuscular blockade, human volunteers, 88:633
- Vasodilation**, see Heart
- Vasopressin**, see Pharmacokinetics; Pharmacology
- Vasopressors**, use after cranial trauma (letter), 88:1423
- Vasovagal episodes**, see Complications
- Vecuronium**, see Neuromuscular relaxants
- Veins**  
cannulation  
external jugular vein, for central venous access (letter), 88:692  
and intracranial pressure, CPB, 88:329  
dorsum vein, on hand, arterialization (letter), 88:1425  
jugular  
external, cannulation, for central venous access (letter), 88:692  
jugular venous oxygen saturation, vs measuring brain tissue oxygenation, for monitoring cerebral oxygenation after traumatic brain injury, 88:549
- Ventilation**  
arterial-endtidal carbon dioxide difference, unaffected during moderate hypothermia, 88:538  
arterio-alveolar oxygen tension difference, increased during moderate hypothermia, 88:538  
depression, sameridine vs morphine, 88:160  
difficulties, through intubating LMA, 88:1181  
end-tidal to arterial CO<sub>2</sub> gradient, and surgical position, 88:43  
function, changes, after interscalene brachial plexus anesthesia with 0.5% and 0.75% ropivacaine, double-blind comparison with 2% mepivacaine, 88:587  
hypoxic ventilatory response, effects of CO<sub>2</sub> and of sustained hypoxia (letter), 88:695  
jet  
high-frequency, for tracheal resection, respiratory gas monitoring using double-lumen jet catheter, 88:224  
in upper airway obstruction, rediscovering the obvious (letter), 88:1427  
mechanical  
dynamic compliance and inspiratory flow, relationship (letter), 88:465  
fiberoptic-guided airway exchange of esophageal-tracheal Combitube, 88:193  
minute, adult circle breathing system vs Bain system, infant lung model with pressure-limited ventilation, 88:527  
one-lung  
arterial oxygenation during, combined vs general anesthesia, 88:494  
reliability of continuous intraarterial blood gas monitoring (letter), 88:1192
- positive end-expiratory pressure, predictors for lung volume reduction surgery, 88:28  
spontaneous, patients with, venous air embolism during Parkinson surgery, 88:793  
volume-controlled, effects of changes in fresh gas flow, Ohmeda 7000 vs Ohmeda 7900 ventilators, 88:39
- Ventilators**, see Equipment
- Video**, preanesthetic, for facilitation of parental education and anxiolysis before pediatric ambulatory surgery, 88:246
- Video imaging**, vs cuff pressure measurement, assessing neuromuscular block at larynx, 88:426
- Visual analog scale**, see Measurement techniques
- Vomiting, nausea and, postoperative (PONV)**  
cesarean section, spinal anesthesia, efficacy of granisetron/dexamethasone combination, 88:1346  
incidence, efficacy and adverse effects of prophylactic antiemetics during PCA, quantitative systematic review, 88:1354  
ondansetron vs droperidol (letter), 88:465  
preventing  
comparative efficacy and safety of ondansetron, droperidol, and metoclopramide, meta-analysis, 88:1370  
use of nonpharmacological techniques, meta-analysis, 88:1362  
TCI vs standard anesthesia regimen, 88:77  
therapies, understanding, and use of meta-analysis (editorial), 88:1200  
tonsillectomy, use of preoperative oral antiemetics, granisetron vs perphenazine, children, 88:1298
- von Hippel-Lindau disease**, see Complications
- Xenon**, see Anesthetics, gases
- Xenopus oocytes**, actions of fluorinated alkanols, 88:877
- Yohimbine**, see Analgesics

